

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 1 113 007 A1**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:

**04.07.2001 Bulletin 2001/27**

(21) Application number: **00311197.8**

(22) Date of filing: **14.12.2000**

(51) Int Cl.<sup>7</sup>: **C07D 217/04**, C07D 217/06,  
C07D 217/14, C07D 217/18,  
C07D 409/04, C07D 401/12,  
C07D 413/12, C07D 405/06,  
C07D 417/04, C07D 401/04,  
A61K 31/472, A61K 31/4725,  
A61P 5/30

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU  
MC NL PT SE TR**

Designated Extension States:

**AL LT LV MK RO SI**

(30) Priority: **24.12.1999 US 173063 P**

(71) Applicant: **PFIZER INC.**

**Groton, CT 06340 (US)**

(72) Inventors:

- **Chesworth, Richard**  
**Groton, Connecticut 06340 (US)**

- **Cameron, Kimberly O'Keefe**  
**Groton, Connecticut 06340 (US)**
- **Da Silva-Jardine, Paul Andrew**  
**Groton, Connecticut 06340 (US)**
- **Day, Robert Francis**  
**Groton, Connecticut 06340 (US)**
- **Lefker, Bruce Allen**  
**Groton, Connecticut 06340 (US)**
- **Zawistoski, Michael Paul**  
**Groton, Connecticut 06340 (US)**

(74) Representative: **Wood, David John et al**  
**PFIZER LIMITED,**  
**Ramsgate Road**  
**Sandwich, Kent CT13 9NJ (GB)**

(54) **Tetrahydroisoquinoline compounds as estrogen agonists/antagonists**

(57) This invention relates to compounds useful for treating or preventing obesity, breast cancer, osteoporosis,

endometriosis, cardiovascular disease, prostatic disease, and the like, and to pharmaceutical composition, methods, and kits comprising such compounds.

**EP 1 113 007 A1**

**Description**

**[0001]** This invention relates to novel tetrahydroisoquinoline compounds that are useful as estrogen agonists and antagonists, and the pharmaceutical uses thereof.

**BACKGROUND OF THE INVENTION**

**[0002]** The value of naturally occurring estrogens and synthetic compositions demonstrating "estrogenic" activity has typically been in their medical and therapeutic uses. A traditional listing of the therapeutic applications for estrogens alone or in combination with other active agents includes, but is not limited to, oral contraception, relief for the symptoms of menopause, prevention of threatened or habitual abortion, relief of dysmenorrhea, relief of dysfunctional uterine bleeding, an aid in ovarian development, treatment of acne, diminution of excessive growth of body hair in women (hirsutism), the prevention of cardiovascular disease, treatment of osteoporosis, treatment of prostatic carcinoma, and suppression of post-partum lactation (Goodman and Gilman, *The Pharmacological Basis Of Therapeutics* (7th Ed.), Macmillan Publishing Company, 1985, pages 1421-1423). Accordingly, there has been increasing interest in finding newly synthesized compounds and new uses for previously known compounds that are demonstrably estrogenic, that is, able to mimic the action of estrogen in estrogen responsive tissue.

**[0003]** From the viewpoint of pharmacologists interested in developing new drugs useful for the treatment of human diseases and specific pathological conditions, it is desirable to procure compounds having demonstrable estrogen-like function, but which are devoid of unwanted side-effects. Exemplifying this latter view, osteoporosis, a disease in which bone becomes increasingly more fragile, is greatly ameliorated by the use of fully active estrogens. However, due to the recognized increased risk of uterine cancer in patients treated chronically with active estrogens, it is not clinically advisable to treat osteoporosis in intact women with fully active estrogens for prolonged periods.

**[0004]** Osteoporosis is a systemic skeletal disease, characterized by low bone mass and deterioration of bone tissue, with a consequent increase in bone fragility and susceptibility to fracture. In the U.S., the condition affects more than 25 million people and causes more than 1.3 million fractures each year, including 500,000 spine, 250,000 hip, and 240,000 wrist fractures annually. These injuries cost the nation over \$10 billion per year. Hip fractures are the most serious, with 5-20% of patients dying within one year, and over 50% of the survivors being incapacitated.

**[0005]** The elderly are at greatest risk of osteoporosis, and the problem is therefore predicted to increase significantly with the aging of the population. Worldwide fracture incidence is forecast to increase 3-fold over the next 60 years, and one study estimates there will be 4.5 million hip fractures worldwide in 2050.

**[0006]** Women are at greater risk of osteoporosis than men. Women experience a sharp acceleration of bone loss during the five years following menopause. Other factors that increase the risk include smoking, alcohol abuse, a sedentary lifestyle, and low calcium intake.

**[0007]** Estrogen is the agent of choice in preventing osteoporosis or post menopausal bone loss in women; it is the only treatment that unequivocally reduces fractures. However, estrogen stimulates the uterus and is associated with an increased risk of endometrial cancer. Although the risk of endometrial cancer is thought to be reduced by a concurrent use of a progestogen, there remains concern about possible increased risk of breast cancer with the use of estrogen.

**[0008]** Black et al., in EP 0605193A1, report that estrogen, particularly when taken orally, lowers plasma levels of LDL and raises plasma levels of the beneficial high density lipoproteins (HDLs). Thus, estrogen can be an effective therapy for hypercholesterolemia. However, as discussed supra, long-term estrogen therapy has been implicated in a variety of disorders, including an increase in the risk of uterine and breast cancer, causing many women to avoid this treatment. Recently suggested therapeutic regimens, that seek to lessen the cancer risk, such as administering combinations of progestogen and estrogen, cause the patient to experience unacceptable bleeding. Furthermore, combining progestogen with estrogen seems to blunt the desired serum cholesterol effects of estrogen. The significant undesirable effects associated with estrogen therapy support the need to develop alternative therapies for hypercholesterolemia that have the desirable effect on serum LDL but do not cause undesirable effects.

**[0009]** There is a need for improved estrogen agonists that exert selective effects on different tissues in the body. Tamoxifen, or 1-(4- $\beta$ -dimethylaminoethoxyphenyl)-1,2-diphenyl-but-1-ene, is an antiestrogen that has a palliative effect on breast cancer, but is reported to have estrogenic activity in the uterus. Gill-Sharma et al., *J. Repr. Fert.*, 99:395 (1993), discloses that tamoxifen at 200 and 400 mg/kg/day reduces the weights of the testes and secondary sex organs in male rats.


**[0010]** Recently it has been reported (Osteoporosis Conf. Scrip No. 1812/13, p. 29 (April 16-20, 1993)) that raloxifene, or 6-hydroxy-2-(4-hydroxyphenyl)-3-[4-(2-piperidinoethoxy) benzoyl] benzo[b] thiophene, mimics the favorable action of estrogen on bone and lipids but, unlike estrogen, has minimal uterine stimulatory effect. (Jordan et al., *Breast Cancer Res. Treat.*, 10(1):31-36 (1987)).

**[0011]** Neubauer et al., *The Prostate*, 23:245 (1993), teaches that raloxifene treatment of male rats produced regression of the ventral prostate.


**[0012]** Raloxifene and related compounds are described as antiestrogenic and antiandrogenic materials that are effective in the treatment of certain mammary and prostate cancers. See U.S. Patent 4,418,068 and Jones et al., *J. Med. Chem.*, 27:1057-66 (1984).

**[0013]** Jones et al. in U.S. Patent 4,133,814 describe derivatives of 2-phenyl-3-aryl-benzothiophene and 2-phenyl-3-arylbenzothiophene-1-oxides that are useful as antifertility agents, and also suppress the growth of mammary tumors.


**[0014]** Lednicer et al., *J. Med. Chem.*, 12:881 (1969), describes estrogen antagonists of the structure



wherein R<sup>2</sup> is phenyl or cyclopentyl and R<sup>3</sup> is H, -CH<sub>2</sub>CHOHCH<sub>2</sub>OH, or




**[0015]** Bencze et al., *J. Med. Chem.*, 10:138 (1967), prepared a series of tetrahydronaphthalenes intended to achieve separation of estrogenic, antifertility, and hypocholesterolemic activities, although they were only partially successful in doing so. These structures have the general formula:




wherein R<sup>1</sup> is H or OCH<sub>3</sub>, R<sup>2</sup> is H, OH, OCH<sub>3</sub>, OPO(OC<sub>2</sub>H<sub>5</sub>)<sub>2</sub>, OCH<sub>2</sub>CH<sub>2</sub>N(C<sub>2</sub>H<sub>5</sub>)<sub>2</sub>, OCH<sub>2</sub>COOH, or OCH(CH<sub>3</sub>)COOH, and R<sup>3</sup> is H or Cl.

**[0016]** U.S. Patent 3,234,090 discloses compounds having estrogenic and antifungal properties, as well as procedures for the preparation of these compounds. The described compounds have the formula:




in which Ph is a 1,2-phenylene radical, Ar is a monocyclic carbocyclic aryl group substituted by tertiary amino-lower alkyl-oxy, in which the tertiary amino is separated from the oxy by at least two carbon atoms, R is hydrogen, an aliphatic radical, a carbocyclic aryl radical, a carbocyclic aryl-aliphatic radical, a heterocyclic aryl radical, or a heterocyclic aryl aliphatic radical, the group of the formula  $-(C_nH_{2n-2})-$  stands for an unbranched alkylene radical having from three to five carbon atoms and carrying the groups Ar and R, salts, N-oxides, salts of N-oxides, or quaternary ammonium compounds thereof.

**[0017]** U.S. Patent 3,277,106 refers to basic ethers with estrogenic, hypocholesterolemic, and antifertility effects, those ethers having the formula:



in which Ph is a 1,2-phenylene radical, Ar is a monocyclic aryl radical substituted by at least one amino-lower alkyl-oxy group in which the nitrogen atom is separated from the oxygen atom by at least two carbon atoms, R is an aryl radical, and the portion  $-(C_nH_{2n-2})-$  stands for lower alkylene forming with Ph a six- or seven-membered ring, two of the ring carbon atoms thereof carry the groups Ar and R, salts, N-oxides, salts of N-oxides, and quaternary ammonium compounds thereof.

**[0018]** Lednicer et al., in *J. Med. Chem.*, 10:78 (1967), and in U.S. Patent 3,274,213, refer to compounds of the formula



wherein  $R_1$  and  $R_2$  are selected from the class consisting of lower alkyl and lower alkyl linked together to form a 5 to 7 ring member saturated heterocyclic radical.

**[0019]** PCT publication No. WO 96/09040 A1 discloses a benzofuran compound useful for treatment of medical indications associated with post-menopausal syndrome, e.g., uterine fibroid disease, endometriosis, and aortal smooth muscle cell proliferation.

**[0020]** European Patent Application EP 0,826,670 A1 discloses naphthalene compounds and methods for inhibiting estrogen deficient pathologies such as lack of birth control, postmenopausal syndrome including osteoporosis, cardiovascular disease, restenosis and hyperlipidemia, prostate cancer, acne, hirsutism, dysfunctional uterine bleeding, dysmenorrhea, and atrophic vaginitis.

**[0021]** European Patent Application EP 0,659,424 A1 discloses benzothiophene compounds useful for treating male infertility.

**[0022]** U.S. Patent 5,462,950 discloses the use of benzothiophene compounds for treating physical menstrual symptoms.

**[0023]** PCT Publication WO 96/40134 discloses methods of antagonizing or blocking calcium channels in vascular tissue comprising administering a benzothiophene compound.

**[0024]** U.S. Patent 5,521,214 discloses methods of inhibiting environmental estrogens comprising administering a benzothiophene compound.

**[0025]** U.S. Patent 5,554,628 discloses methods for minimizing the uterotrophic effect of tamoxifen and its analogs by administering a naphthalene compound.

**[0026]** PCT Publication WO 97/13764 discloses benzothiophene compounds useful for inhibiting cardiovascular disease including restenosis and atherosclerosis.

**[0027]** PCT Publication WO 97/13511 discloses benzothiophene compounds useful for inhibiting plasminogen activator inhibitor 1 related conditions such as major tissue damage and trauma, or multiple organ dysfunction syndrome.

**[0028]** European Patent Application 0,729,754 A2 discloses benzothiophene compounds useful for inhibiting estrogen positive tumors of the brain or CNS.

**[0029]** U.S. Patent 5,670,523 discloses benzothiophene compounds useful for inhibiting musculoaponeurotic fibromatoses previously classified as desmoid tumors.

**[0030]** U.S. Patent 5,496,828 discloses methods for inhibiting ulcerative mucositis by administering a benzothio-  
 phene compound.

**[0031]** PCT Publication WO 97/26876 discloses methods for increasing sphincter competence by administering a benzothiophene compound.

**[0032]** European Patent Application 0,826,679 A2 discloses a pharmaceutical composition useful for alleviating symptoms of postmenopausal syndromes, the composition comprising a naphthalene compound and an additional therapeutic agent such as estrogen or progestin, a benzothiophene compound such as raloxifene, a naphthyl compound having antiestrogen activity, a bisphosphonate compound such as alendronate or tiludronate, parathyroid hormone (PTH), including truncated and/or recombinant forms of PTH such as PTH (1-34), calcitonin, bone morphogenic proteins (BMPs), or combination thereof.


**[0033]** European Patent Application 0,702,962 A2 discloses a pharmaceutical agent for treating breast cancer, the agent comprising tamoxifen and a naphthyl compound useful for inhibiting hormone-dependent breast cancer.

**[0034]** U.S. Patent 5,599,822 discloses a pharmaceutical composition for minimizing the bone loss effect of danazol, the composition comprising danazol and a benzothiophene compound having antiestrogen activity.

## SUMMARY OF THE INVENTION

**[0035]** This invention relates to novel tetrahydroisoquinoline compounds that are useful as estrogen agonists and antagonists, and the pharmaceutical uses thereof.

[0036] In a first aspect, this invention provides compounds of the formula:



wherein:

A<sup>1</sup> is hydrogen, hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, or (C<sub>1</sub>-C<sub>4</sub>)alkanoyloxy, said (C<sub>1</sub>-C<sub>4</sub>)alkoxy or said (C<sub>1</sub>-C<sub>4</sub>)alkanoyloxy being optionally substituted by hydroxy, halo, or a partially saturated, fully saturated, or fully unsaturated five to twelve membered ring optionally having up to four heteroatoms independently selected from oxygen, sulfur, and nitrogen, or A<sup>1</sup> is R<sup>3</sup>-(C<sub>1</sub>-C<sub>4</sub>)alkoxy wherein R<sup>3</sup> is pyrrolidino, piperidino, morpholino, or dimethylamino;

A<sup>2</sup>, A<sup>3</sup>, and A<sup>4</sup> are independently selected from hydrogen, hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, and halo;

R<sup>1</sup> is phenyl; pyridyl; piperidinyl; (C<sub>1</sub>-C<sub>7</sub>)alkyl; adamantyl; a partially saturated, fully saturated, or fully unsaturated three to twelve membered ring optionally having up to four heteroatoms selected independently from oxygen, sulfur, and nitrogen; a bicyclic ring consisting of two fused independently partially saturated, fully saturated, or fully unsaturated five to six membered rings, wherein said bicyclic ring includes up to four heteroatoms independently

selected from oxygen, sulfur, and nitrogen; or abicyclic ring system consisting of two rings joined by a covalent bond, said rings being independently partially saturated, fully saturated, or fully unsaturated three to eight membered rings, wherein said bicyclic ring system includes up to four heteroatoms independently selected from oxygen, sulfur, and nitrogen; wherein each of the above R<sup>1</sup> groups is optionally substituted with up to seven fluoro atoms, or with up to three substituents independently selected from Group A, wherein Group A consists of hydroxy, halo, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, R<sup>3</sup>-(C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>2</sub>-C<sub>4</sub>)alkenyl-COOR<sup>7</sup> wherein R<sup>7</sup> is hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-COOR<sup>7</sup>, (C<sub>1</sub>-C<sub>4</sub>)alkanoyloxy-(C<sub>2</sub>-C<sub>4</sub>)alkenyl, (C<sub>2</sub>-C<sub>4</sub>)alkenyl-CONR<sup>4</sup>R<sup>5</sup> wherein R<sup>4</sup> and R<sup>5</sup> are independently hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkylene, or (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, or R<sup>4</sup> and R<sup>5</sup> taken together with the nitrogen atom to which they are attached form pyrrolidino, piperidino, morpholino, or hexamethyleneimino, (C<sub>0</sub>-C<sub>4</sub>)alkyl-CONR<sup>4</sup>R<sup>5</sup>, (C<sub>0</sub>-C<sub>4</sub>)alkyl-NR<sup>4</sup>R<sup>5</sup>, OCH<sub>2</sub>CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup> wherein R<sup>8</sup> and R<sup>9</sup> are independently methyl or ethyl, or R<sup>8</sup> and R<sup>9</sup> taken together with the nitrogen atom to which they are attached form pyrrolidino, piperidino, morpholino, or hexamethyleneimino, propyl-R<sup>8</sup>R<sup>9</sup>, and SO<sub>2</sub>-R<sup>6</sup> wherein R<sup>6</sup> is imidazolyl, thienyl, benzathienyl, or isoxazolyl, optionally substituted with up to three substituents independently selected from (C<sub>1</sub>-C<sub>4</sub>)alkyl;

X is a covalent bond, (CH<sub>2</sub>)<sub>n</sub> where n is 1, 2, or 3, (C<sub>0</sub>-C<sub>1</sub>)alkylene-phenylene-(C<sub>0</sub>-C<sub>1</sub>)alkylene, CO<sub>2</sub>, (C<sub>0</sub>-C<sub>3</sub>)alkylene-CO-(C<sub>0</sub>-C<sub>3</sub>)alkylene, or (C<sub>0</sub>-C<sub>4</sub>)alkylene-SO<sub>2</sub>-(C<sub>0</sub>-C<sub>4</sub>)alkylene;

R<sup>2</sup> is (C<sub>1</sub>-C<sub>9</sub>)alkyl; (C<sub>2</sub>-C<sub>4</sub>)alkenyl; benzhydryl; a partially saturated, fully saturated, or fully unsaturated three to eight membered ring optionally having up to four heteroatoms selected independently from oxygen, sulfur, and nitrogen; a bicyclic ring consisting of two fused independently partially saturated, fully saturated, or fully unsaturated five to six membered rings, wherein said bicyclic ring includes up to four heteroatoms independently selected from oxygen, sulfur, and nitrogen; or a bicyclic ring system consisting of two rings joined by a covalent bond, said rings being independently partially saturated, fully saturated, or fully unsaturated three to eight membered rings, wherein said bicyclic ring system includes up to four heteroatoms independently selected from oxygen, sulfur, and nitrogen; wherein said (C<sub>1</sub>-C<sub>9</sub>)alkyl is optionally substituted with one to seven fluoro substituents, or up to three substituents independently selected from Group B, wherein Group B consists of chloro, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, amino, and (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonyl; wherein said (C<sub>2</sub>-C<sub>4</sub>)alkenyl is optionally substituted with up to three substituents independently selected from Group C, wherein Group C consists of halo, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, amino, and (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonyl; and wherein said benzhydryl, said 5 to 8 membered ring, said bicyclic ring, and said bicyclic ring system is optionally substituted with up to three substituents independently selected from Group D, wherein Group D consists of halo, hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, imidazolyl, amino, (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonylamino, and (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonyl; and p is 0, 1, or 2;

with the proviso that

when X is (CH<sub>2</sub>)<sub>2</sub> or (CH<sub>2</sub>)<sub>3</sub>, p is 0, and R<sup>1</sup> is phenyl or phenyl substituted with a single chloro, fluoro, bromo, hydroxy, methoxy, pyrrolidinoethoxy, piperidinoethoxy, or morpholinoethoxy substituent, then R<sup>2</sup> is not phenyl, methoxyphenyl, tert-butyl, or cyclopentyl;  
 when X is CH<sub>2</sub>, (CH<sub>2</sub>)<sub>2</sub>, COCH<sub>2</sub>, or CH<sub>2</sub>CO, A<sup>1</sup> is hydrogen, and R<sup>1</sup> is phenyl, then R<sup>2</sup> is not phenyl; and  
 when X is a covalent bond, p is 0, A<sup>1</sup> is hydrogen or methoxy, and R<sup>1</sup> is phenyl or phenyl substituted with a single chloro, fluoro, bromo, methoxy, pyrrolidinoethoxy, or piperidinoethoxy substituent, then R<sup>2</sup> is not phenyl or *m*-fluorophenyl.

[0037] In a preferred embodiment of the first aspect, A<sup>1</sup> is hydroxy; A<sup>2</sup>, A<sup>3</sup>, and A<sup>4</sup> are hydrogen; and p is 0.

[0038] In another preferred embodiment of the first aspect, R<sup>1</sup> is phenyl, pyridyl, (C<sub>1</sub>-C<sub>4</sub>)alkyl, adamantyl, naphthyl, or a partially saturated, fully saturated, or fully unsaturated five to six membered ring optionally having up to two heteroatoms selected independently from oxygen, sulfur, and nitrogen; wherein each of said R<sup>1</sup> groups is optionally substituted with up to seven fluoro atoms, or with up to three substituents independently selected from Group A.

[0039] In another preferred embodiment of the first aspect, R<sup>1</sup> is phenyl, cyclohexyl, pyridyl, thienyl, isopropyl, or adamantyl; wherein each of said R<sup>1</sup> groups is optionally substituted with up to seven fluoro atoms, or with up to three substituents independently selected from Group A.

[0040] In another preferred embodiment of the first aspect, R<sup>1</sup> is phenyl or cyclohexyl; wherein each of said R<sup>1</sup> groups is optionally substituted with up to seven fluoro atoms, or with up to three substituents independently selected from Group A.

[0041] In another preferred embodiment of the first aspect, each of said R<sup>1</sup> groups is substituted with up to three halo atoms, or with one substituent selected from hydroxy, (C<sub>1</sub>-C<sub>2</sub>)alkoxy, pyrrolidino-(C<sub>1</sub>-C<sub>4</sub>)alkoxy, dimethylamino, (C<sub>2</sub>-C<sub>4</sub>)alkenyl-COOR<sup>7</sup>, COOR<sup>7</sup>, (C<sub>2</sub>-C<sub>4</sub>)alkenyl-CONR<sup>4</sup>R<sup>5</sup> wherein R<sup>4</sup> and R<sup>5</sup> are independently hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkyl, -(CH<sub>2</sub>CH<sub>2</sub>-O-CH<sub>3</sub>), or (C<sub>5</sub>-C<sub>6</sub>)cycloalkyl, or R<sup>4</sup> and R<sup>5</sup> taken together with the nitrogen atom to which they are attached form piperidino or morpholino, or SO<sub>2</sub>-R<sup>6</sup> wherein R<sup>6</sup> is imidazolyl optionally substituted

with up to three substituents independently selected from (C<sub>1</sub>-C<sub>4</sub>)alkyl.

**[0042]** In another preferred embodiment of the first aspect, each of said R<sup>1</sup> groups is substituted with up to three fluoro atoms, or with one substituent selected from iodo, chloro, bromo, hydroxy, methoxy, pyrrolidino-ethoxy, dimethylamino, COOR<sup>7</sup> wherein R<sup>7</sup> is hydrogen or methyl, or ethenyl-CONR<sup>4</sup>R<sup>5</sup> wherein R<sup>4</sup> and R<sup>5</sup> are both methyl, or R<sup>4</sup> and R<sup>5</sup> taken together with the nitrogen atom to which they are attached form piperidino or morpholino.

**[0043]** In another preferred embodiment of the first aspect, each of said R<sup>1</sup> groups is substituted with one hydroxy or pyrrolidino-ethoxy.

**[0044]** In another preferred embodiment of the first aspect, X is a covalent bond, CH<sub>2</sub>, CH<sub>2</sub>-phenylene, CO<sub>2</sub>, CO-(C<sub>0</sub>-C<sub>2</sub>)alkylene, or SO<sub>2</sub>-(C<sub>0</sub>-C<sub>2</sub>)alkylene.

**[0045]** In another preferred embodiment of the first aspect, X is a covalent bond, CO, or SO<sub>2</sub>.

**[0046]** In another preferred embodiment of the first aspect, R<sup>2</sup> is (C<sub>1</sub>-C<sub>7</sub>)alkyl; propenyl; a partially saturated, fully saturated, or fully unsaturated five to seven membered ring optionally having up to two heteroatoms selected independently from oxygen, sulfur, and nitrogen; a bicyclic ring consisting of two fused independently partially saturated, fully saturated, or fully unsaturated five to six membered rings, wherein said bicyclic ring includes up to two oxygen atoms; or biphenyl; wherein said (C<sub>1</sub>-C<sub>7</sub>)alkyl is optionally substituted with one to seven fluoro substituents, or up to three substituents independently selected from Group B; wherein said propenyl is optionally substituted with up to three substituents independently selected from Group C; and wherein each of said 5-7 membered ring, said bicyclic ring, and said biphenyl is optionally substituted with up to three substituents independently selected from Group D. In an even more preferred embodiment, each of said (C<sub>1</sub>-C<sub>7</sub>)alkyl and said propenyl is substituted with one to three fluoro substituents, or up to two substituents independently selected from amino and methylcarbonyl; and wherein each of said 5-7 membered ring, said bicyclic ring, and said biphenyl is substituted with up to three fluoro substituents, or up to two substituents independently selected from hydroxy, (C<sub>1</sub>-C<sub>3</sub>)alkyl, amino, and methylcarbonyl.

**[0047]** In another preferred embodiment of the first aspect, R<sup>2</sup> is methyl, t-butyl, phenyl, cyclohexyl, isoxazolyl, tetrahydropyranyl, naphthyl, or benzodioxolyl; wherein each of said methyl or t-butyl is optionally substituted with one to seven fluoro substituents, or up to three substituents independently selected from Group B; and wherein each of said phenyl, cyclohexyl, isoxazolyl, tetrahydropyranyl, naphthyl, or benzodioxolyl is optionally substituted with up to three substituents independently selected from Group D.

**[0048]** In another preferred embodiment of the first aspect, R<sup>2</sup> is trifluoromethyl or phenyl; wherein said phenyl is optionally substituted with up to three substituents independently selected from Group D.

**[0049]** In another preferred embodiment of the first aspect,

A<sup>1</sup> is hydroxy;

A<sup>2</sup>, A<sup>3</sup>, and A<sup>4</sup> are hydrogen;

p is 0;

R<sup>1</sup> is phenyl, cyclohexyl, pyridyl, thienyl, isopropyl, or adamantyl; wherein each of said R<sup>1</sup> groups is optionally substituted with up to three fluoro atoms, or with one substituent selected from iodo, chloro, bromo, hydroxy, methoxy, pyrrolidino-ethoxy, dimethylamino, COOR<sup>7</sup> wherein R<sup>7</sup> is hydrogen or methyl, or ethenyl-CONR<sup>4</sup>R<sup>5</sup> wherein R<sup>4</sup> and R<sup>5</sup> are both methyl, or R<sup>4</sup> and R<sup>5</sup> taken together with the nitrogen atom to which they are attached form piperidino or morpholino;

X is a covalent bond, CH<sub>2</sub>, CH<sub>2</sub>-phenylene, CO<sub>2</sub>, CO-(C<sub>0</sub>-C<sub>2</sub>)alkylene, or SO<sub>2</sub>-(C<sub>0</sub>-C<sub>2</sub>)alkylene; and

R<sup>2</sup> is methyl, t-butyl, phenyl, cyclohexyl, isoxazolyl, tetrahydropyranyl, naphthyl, or benzodioxolyl; wherein each of said methyl or t-butyl is optionally substituted with one to three fluoro substituents, or up to two substituents independently selected from amino and methylcarbonyl; and wherein each of said phenyl, cyclohexyl, isoxazolyl, tetrahydropyranyl, naphthyl, or benzodioxolyl is optionally substituted with up to three fluoro substituents, or up to two substituents independently selected from hydroxy, (C<sub>1</sub>-C<sub>3</sub>)alkyl, amino, and methylcarbonyl.

**[0050]** Some of the more preferred compounds of the first aspect of the present invention include 1-(4-hydroxyphenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinolin-6-ol; 3-[4-(6-hydroxy-2-phenyl-1,2,3,4-tetrahydroisoquinolin-1-yl)-phenyl]-1-piperidin-1-yl-propenone; 3-[4-(6-hydroxy-2-phenyl-1,2,3,4-tetrahydroisoquinolin-1-yl)-phenyl]-1-morpholin-4-yl-propenone; 3-[4-(6-hydroxy-2-phenyl-1,2,3,4-tetrahydroisoquinolin-1-yl)-phenyl]-N,N-dimethyl-acrylamide; 2-benzyl-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol; 2,2,2-trifluoro-1-[6-hydroxy-1-(4-hydroxyphenyl)-3,4-dihydro-1H-isoquinolin-2-yl]-ethanone; 2-benzenesulfonyl-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol; and 2-(4-isopropylbenzenesulfonyl)-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol.

**[0051]** In another preferred embodiment of the first aspect,

said compound is of formula (I);

A<sup>1</sup> is hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>1</sub>-C<sub>4</sub>)alkanoyloxy, or pyrrolidino-ethoxy;





A<sup>2</sup>, A<sup>3</sup>, and A<sup>4</sup> are independently selected from hydrogen, hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, and halo;

R<sup>1</sup> is phenyl; pyridyl; piperidyl; (C<sub>1</sub>-C<sub>7</sub>)alkyl; adamantyl; a partially saturated, fully saturated, or fully unsaturated three to twelve membered ring optionally having up to four heteroatoms selected independently from oxygen, sulfur, and nitrogen; a bicyclic ring consisting of two fused independently partially saturated, fully saturated, or fully unsaturated five to six membered rings, wherein said bicyclic ring includes up to four heteroatoms independently selected from oxygen, sulfur, and nitrogen; or abicyclic ring system consisting of two rings joined by a covalent bond, said rings being independently partially saturated, fully saturated, or fully unsaturated three to eight membered rings, wherein said bicyclic ring system includes up to four heteroatoms independently selected from oxygen, sulfur, and nitrogen; wherein each of the above R<sup>1</sup> groups is optionally substituted with up to seven fluoro atoms, or with up to three substituents independently selected from Group A, wherein Group A consists of hydroxy, halo, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, R<sup>3</sup>-(C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>2</sub>-C<sub>4</sub>)alkenyl-COOR<sup>7</sup> wherein R<sup>7</sup> is hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-COOR<sup>7</sup>, (C<sub>1</sub>-C<sub>4</sub>)alkanoyloxy-(C<sub>2</sub>-C<sub>4</sub>)alkenyl, (C<sub>2</sub>-C<sub>4</sub>)alkenyl-CONR<sup>4</sup>R<sup>5</sup> wherein R<sup>4</sup> and R<sup>5</sup> are independently hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkylene, or (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, or R<sup>4</sup> and R<sup>5</sup> taken together with the nitrogen atom to which they are attached form pyrrolidino, piperidino, morpholino, or hexamethyleneimino, (C<sub>0</sub>-C<sub>4</sub>)alkyl-CONR<sup>4</sup>R<sup>5</sup>, (C<sub>0</sub>-C<sub>4</sub>)alkyl-NR<sup>4</sup>R<sup>5</sup>, OCH<sub>2</sub>CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup> wherein R<sup>8</sup> and R<sup>9</sup> are independently methyl or ethyl, or R<sup>8</sup> and R<sup>9</sup> taken together with the nitrogen atom to which they are attached form pyrrolidino, piperidino, morpholino, or hexamethyleneimino, propyl-R<sup>8</sup>R<sup>9</sup>, and SO<sub>2</sub>-R<sup>6</sup> wherein R<sup>6</sup> is imidazolyl, thienyl, benzathienyl, or isoxazolyl, optionally substituted with up to three substituents independently selected from (C<sub>1</sub>-C<sub>4</sub>)alkyl;

X is (C<sub>0</sub>-C<sub>1</sub>)alkylene-phenylene-(C<sub>0</sub>-C<sub>1</sub>)alkylene, CO<sub>2</sub>, CO, (C<sub>1</sub>-C<sub>3</sub>)alkylene-CO-(C<sub>1</sub>-C<sub>3</sub>)alkylene, (C<sub>0</sub>-C<sub>3</sub>)alkylene-CO-(C<sub>2</sub>-C<sub>3</sub>)alkylene, (C<sub>2</sub>-C<sub>3</sub>)alkylene-CO-(C<sub>0</sub>-C<sub>3</sub>)alkylene, or (C<sub>0</sub>-C<sub>4</sub>)alkylene-SO<sub>2</sub>-(C<sub>0</sub>-C<sub>4</sub>)alkylene; R<sup>2</sup> is (C<sub>1</sub>-C<sub>9</sub>)alkyl; (C<sub>2</sub>-C<sub>4</sub>)alkenyl; benzhydryl; a partially saturated, fully saturated, or fully unsaturated three to eight membered ring optionally having up to four heteroatoms selected independently from oxygen, sulfur, and nitrogen; a bicyclic ring consisting of two fused independently partially saturated, fully saturated, or fully unsaturated five to six membered rings, wherein said bicyclic ring includes up to four heteroatoms independently selected from oxygen, sulfur, and nitrogen; or a bicyclic ring system consisting of two rings joined by a covalent bond, said rings being independently partially saturated, fully saturated, or fully unsaturated three to eight membered rings, wherein said bicyclic ring system includes up to four heteroatoms independently selected from oxygen, sulfur, and nitrogen; wherein said (C<sub>1</sub>-C<sub>9</sub>)alkyl is optionally substituted with one to seven fluoro substituents, or up to three substituents independently selected from Group B, wherein Group B consists of chloro, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, amino, and (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonyl; wherein said (C<sub>2</sub>-C<sub>4</sub>)alkenyl is optionally substituted with up to three substituents independently selected from Group C, wherein Group C consists of halo, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, amino, and (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonyl; and wherein said benzhydryl, said 5 to 8 membered ring, said bicyclic ring, and said bicyclic ring system is optionally substituted with up to three substituents independently selected from Group D, wherein Group D consists of halo, hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, imidazolyl, amino, (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonylamino, and (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonyl; and p is 0, 1, or 2.

**[0056]** In a third aspect, this invention provides methods for treating or preventing a disease, disorder, condition, or symptom mediated by an estrogen receptor and/or caused by lowered estrogen level in a mammal, said method comprising administering to said mammal a therapeutically effective amount of a compound of the present invention.

**[0057]** In preferred embodiments of the third aspect, a therapeutically effective amount of a compound of the present invention is combined with a therapeutically effective amount of an anabolic agent, a prodrug thereof, or a pharmaceutically acceptable salt of said anabolic agent or said prodrug; a growth hormone or a growth hormone secretagogue, a prodrug thereof, or a pharmaceutically acceptable salt of said growth hormone secretagogue or said prodrug; a prostaglandin agonist/antagonist, a prodrug thereof, or a pharmaceutically acceptable salt of said prostaglandin agonist/antagonist or said prodrug; or a parathyroid hormone or sodium fluoride.

**[0058]** In another preferred embodiment of the third aspect, said disease, disorder, condition, or symptom is perimenopausal or postmenopausal syndrome, osteoporosis, atrophy of skin or vagina, elevated serum cholesterol levels, cardiovascular disease, Alzheimer's disease, a reduction or prevention of reduction in cognitive function, an estrogen dependent cancer, breast or uterus cancer, a prostatic disease, benign prostatic hyperplasia, or prostate cancer.

**[0059]** In another preferred embodiment of the third aspect, said disease, disorder, condition, or symptom is obesity, endometriosis, bone loss, uterine fibrosis, aortal smooth muscle cell proliferation, lack of birth control, acne, hirsutism, dysfunctional uterine bleeding, dysmenorrhea, male infertility, impotence, psychological and behavioral symptoms during menstruation, ulcerative mucositis, uterine fibroid disease, restenosis, atherosclerosis, musculoaponeurotic fibromatosis, alopecia, wound-healing, scarring, auto immune disease, cartilage degeneration, delayed puberty, demyelinating disease, dysmyelinating disease, hypoglycemia, lupus erythematosus, myocardial infarction, ischemia, thromboembolic disorder, obsessive compulsive disorder, ovarian dysgenesis, post menopausal CNS disorder, pulmonary hypertension, reperfusion damage, resistant neoplasm, rheumatoid arthritis, seborrhea, sexual precocity, thyroiditis,

Turner's syndrome, or hyperlipidemia.

**[0060]** In another preferred embodiment of the third aspect, said method is useful for blocking a calcium channel, inhibiting an environmental estrogen, minimizing the uterotrophic effect of tamoxifen or an analog thereof, removing fibrin by inhibiting plasminogen activators, inhibiting estrogen positive primary tumors of the brain and CNS, increasing sphincter competence, increasing libido, inhibiting fertility, oxidizing low density lipoprotein, increasing macrophage function, expressing thrombomodulin, or increasing levels of endogenous growth hormone.

**[0061]** In a fourth aspect, this invention provides pharmaceutical compositions comprising a compound of the present invention and a pharmaceutically acceptable vehicle, carrier, or diluent.

**[0062]** In a fifth aspect, this invention provides kits useful for treating or preventing a disease, disorder, condition, or symptom mediated by an estrogen receptor and/or caused by lowered estrogen levels, said kit comprising a compound of the present invention and a pharmaceutically acceptable vehicle, carrier, or diluent in a dosage form, and a container for containing said dosage form.

**[0063]** In a preferred embodiment of the fifth aspect, said kit also comprises an anabolic agent, a prodrug thereof, or a pharmaceutically acceptable salt of said anabolic agent or said prodrug; a growth hormone or a growth hormone secretagogue, a prodrug thereof, or a pharmaceutically acceptable salt of said growth hormone secretagogue or said prodrug; a prostaglandin agonist/antagonist, a prodrug thereof, or a pharmaceutically acceptable salt of said prostaglandin agonist/antagonist or said prodrug; or a parathyroid hormone or sodium fluoride.

#### DETAILED DESCRIPTION OF THE INVENTION

**[0064]** Halo or halogen refers to chloro, bromo, iodo and fluoro.

**[0065]** Estrogen agonists are herein defined as chemical compounds capable of binding to estrogen receptor sites in mammalian tissue and mimicking the action(s) of estrogen in one or more tissues.

**[0066]** Estrogen antagonists are herein defined as chemical compounds capable of binding to estrogen receptor sites in mammalian tissue and blocking the action(s) of estrogen or estrogen agonists in one or more tissues.

**[0067]** As used in this application, prostatic disease means benign prostatic hyperplasia or prostatic carcinoma.

**[0068]** A "compound", when used to refer to compounds of the present invention, includes within its scope not just the specific compound(s) listed or described, but also alternative forms of the compound, such as prodrugs, pharmaceutically acceptable salts, pharmaceutically acceptable salts of the prodrug, solvates, hydrates, and the like.

**[0069]** A prodrug is a chemical compound that, in its present form, has little or reduced pharmaceutical activity, but which, upon introduction into its biological environment, is readily converted into an active drug form. An exemplary prodrug is an ester of a drug, where upon introduction into a patient, the ester is cleaved to produce the active drug.

**[0070]** A "therapeutically effective amount" of a compound is an amount that is sufficient to cure, prevent, or alleviate a disease, disorder, condition, or symptom.

**[0071]** An "anabolic agent" is any compound that is capable of promoting synthetic metabolic reactions in a patient. In a preferred embodiment, anabolic agents are useful in promoting wound healing.

**[0072]** A "growth hormone or growth hormone secretagogue" is a naturally occurring growth hormone as understood by those skilled in the art, a compound that mimics one or more actions of a naturally occurring growth hormone, or a compound that stimulates the release of naturally occurring growth hormone in a patient.

**[0073]** A "prostaglandin agonist or antagonist" is any compound that is capable of agonizing or antagonizing the activity of a prostaglandin in a patient.

**[0074]** Those of skill in the art will recognize that certain substituents listed in this invention will be chemically incompatible with one another or with the heteroatoms in the compounds, and will avoid these incompatibilities in selecting compounds of this invention. Likewise, certain functional groups may require protecting groups during synthetic procedures as will be recognized by those of skill in the art.

**[0075]** Those of skill in the art will also recognize that certain compounds of this invention will consist of multiple possible geometric configurations, or isomers. All such isomers are included in this invention.

**[0076]** The pharmaceutical compositions and methods of this invention comprise, as an active ingredient, a compound of formula I or II. The pharmaceutically acceptable salts of the compounds of formula I and II are salts of non-toxic type commonly used, such as salts with organic acids (e.g., formic, acetic, trifluoroacetic, citric, maleic, tartaric, methanesulfonic, benzenesulfonic, toluenesulfonic acids, and the like), inorganic acids (e.g. hydrochloric, hydrobromic, sulfuric, phosphoric acids, and the like), amino acids (e.g., aspartic acid, glutamic acid, and the like), alkali metals (e.g., sodium, potassium, and the like), and alkaline earth metals (e.g., calcium, magnesium, and the like). These salts may be prepared by methods known to those of skill in the art.

**[0077]** The compounds of this invention may be administered to animals (including humans) orally, parenterally, topically, or otherwise, in any conventional form of preparation, such as capsules, microcapsules, tablets, granules, powder, troches, pills, suppositories, injections, suspensions, syrups, and the like.

**[0078]** The pharmaceutical compositions of this invention can be prepared by methods commonly employed using

conventional additives, such as excipients (e.g., sucrose, starch, mannitol, sorbitol, lactose, glucose, cellulose, talc, calcium phosphate, calcium carbonate, and the like), binders (e.g., cellulose, methylcellulose, hydroxymethylcellulose, polypropylpyrrolidone, polyvinylpyrrolidone, gelatin, gum arabic, polyethyleneglycol, sucrose, starch, and the like), disintegrators (e.g., starch, carboxymethylcellulose, hydroxypropylstarch, low substituted hydroxypropylcellulose, sodium bicarbonate, calcium phosphate, calcium citrate, and the like), lubricants (e.g., magnesium stearate, light anhydrous silicic acid, talc, sodium lauryl sulfate, and the like), flavoring agents (e.g., citric acid, menthol, glycine, orange powder, and the like), preservatives (e.g., sodium benzoate, sodium bisulfite, methylparaben, propylparaben, and the like), stabilizers (e.g., citric acid, sodium citrate, acetic acid, and the like), suspending agents (e.g., methylcellulose, polyvinylpyrrolidone, aluminum stearate, and the like), dispersing agents (e.g., hydroxypropylmethylcellulose and the like), diluents (e.g., water, alcohol, glycerin, and the like), and base waxes (e.g., cocoa butter, white petrolatum, polyethylene glycol, and the like).

**[0079]** The compounds of this invention may be administered once a day or in multiple daily doses, with a preferred daily dosage of about 0.001 to about 100 mg in adult human patients. This dosage may be properly varied depending on the age, body weight, and medical condition of the patient, as well as the mode of administration. A more preferred daily dose is about 1.0 to about 10 mg in human patients. One dose per day is preferred. Controlled release, sustained release, and/or delayed release oral or parenteral compositions may be used.


#### General Reaction Schemes:

**[0080]** Compounds of this invention are readily prepared by the reactions illustrated in the schemes below.

**[0081]** If required in the following processes, active groups may be protected with a suitable protecting group such as benzyl, p-toluenesulfonyl, methyl, p-methoxybenzyl, and the like, and the protecting group may subsequently be removed at a later stage in the synthesis. The protection and removal of the protecting groups may be carried out according to procedures known to those skilled in the art (e.g., procedures disclosed in *Protective Groups in Organic Synthesis* by T. W. Greene, published by John Wiley & Sons (1991)).

#### General Reaction Scheme A

**[0082]** A compound of formula (Ia) (i.e., a compound of formula (I) wherein X is CO- (CH<sub>2</sub>)<sub>y</sub>, and y is 0-3) may be prepared according to the procedures illustrated in Scheme A.



Scheme A

**[0083]** As illustrated in Scheme A, a phenethyl amine of formula (II) is coupled with a carboxylic acid of formula (III) to give the benzamide of formula (V). Compound (V) is subjected to cyclodehydration to give the dihydroisoquinoline of formula (VI). Dihydroisoquinoline (VI) is reduced to the tetrahydroisoquinoline of formula (VII). Then, compound (VII) may be reacted with a desired acid anhydride, acid chloride, or other coupling agent to give compound (Ia). Alternatively, compound (VII) may be reacted with a desired acid in the presence of 1-propanephosphonic acid cyclic anhydride,  $\text{Et}_3\text{N}$ , and DMAP in dichloromethane ( $\text{CH}_2\text{Cl}_2$ ) to yield compound (Ia).

**[0084]** The coupling of compounds (II) and (III) is carried out in the presence of a coupling agent for peptide synthesis, preferably with an additive. Suitable coupling agents include water-soluble diimides such as N-cyclohexyl-N'-(4-diethylaminocyclohexyl)-carbodiimide and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (EDC), and suitable additives include 1-hydroxybenzotriazole (HOBt) and 3-hydroxy-4-oxo-3,4-dihydro-1,2,3-benzotriazine. This reaction mixture is preferably stirred at a temperature between about  $0^\circ\text{C}$  to room temperature (i.e., about  $25^\circ\text{C}$ ), for about 12 to 60 hours in a reaction inert solvent such as  $\text{CH}_2\text{Cl}_2$ , preferably under nitrogen.

**[0085]** Cyclodehydration of the intermediate compound (V) is carried out in a reaction inert solvent such as 1,2-dichloroethane in the presence of phosphorus pentachloride under nitrogen, or neat using phosphorus oxychloride under nitrogen. Phosphorus pentachloride is added to the solution of compound (V) at about  $0^\circ\text{C}$ , then the reaction mixture is refluxed for about 1 to 24 hours. The reaction of compound (V) and phosphorus oxychloride is carried out at about the reflux temperature of the reaction mixture for about 1 to 48 hours.

**[0086]** The reduction of compound (VI) to compound (VII) is typically carried out in the presence of sodium borohydride in a reaction inert solvent such as methanol or tetrahydrofuran, at from about  $0^\circ\text{C}$  to about room temperature,

for about 1 to 12 hours, preferably under nitrogen. The reaction is quenched with water.

**[0087]** The intermediate compound (VII) thus obtained is reacted with a desired acid anhydride or acid chloride to yield compound (Ia) according to amide formation procedures known to those skilled in the art. This reaction is performed in the presence of a base such as triethylamine ( $\text{Et}_3\text{N}$ ), in a reaction inert solvent such as  $\text{CH}_2\text{Cl}_2$ , for about 1 to 72 hours, at about  $0^\circ\text{C}$  to about room temperature. This reaction is preferably performed under nitrogen.

**[0088]** Alternatively, compound (V) may be prepared by reacting compound (II) with an acyl chloride of formula (IV) in the presence of a base such as triethylamine in a reaction inert solvent. Suitable solvents include dichloromethane, tetrahydrofuran (THF),  $\text{CHCl}_3$ , 1,2-dichloroethane, dioxane, toluene, benzene, and the like, with THF being the most preferred. This reaction is preferably carried out at a temperature of about  $0^\circ\text{C}$  to  $30^\circ\text{C}$ , most preferably about  $25^\circ\text{C}$ , under nitrogen, for about 1 to 24 hours, preferably about 12 hours. If required, a catalyst such as 4-(dimethylamino)pyridine, scandium triflate, tributyl phosphine, or the like may be added to the reaction mixture.

**[0089]** Intermediate compound (VI) may be substituted with a desired substituent on  $\text{R}^1$  and subsequently subjected to the remaining processes illustrated in Scheme A. This substitution reaction is performed according to procedures known to those skilled in the art. For example, compound (VI) wherein  $\text{R}^1$  is piperidin-4-yl may be reacted with a heteroaryl sulfonyl chloride in the presence of a base such as triethylamine in a reaction inert solvent such as dichloroethane to yield a sulfonamide. This reaction is preferably carried out at about room temperature, under nitrogen, for about 1 to 24 hours, preferably about 2 to 20 hours.


**[0090]** A compound of formula (I) of this invention wherein X is  $-(\text{CH}_2)_z-\text{SO}_2-$  may be prepared by reacting compound (VII) with a sulfonyl chloride of formula  $\text{R}^2-(\text{CH}_2)_z-\text{SO}_2-\text{Cl}$  according to known procedures. For example, this reaction may be carried out in the presence of triethylamine in a reaction inert solvent such as tetrahydrofuran (THF) at about room temperature for about 1 to 24 hours under nitrogen.

**[0091]** A compound of formula (I) of this invention wherein X is a covalent bond,  $(\text{CH}_2)_n$ , or  $(\text{CH}_2)_x$ -phenyl (wherein x is 0 or 1), may also be prepared by reacting compound (VII) with an aldehyde of formula  $\text{R}^2-\text{X}-\text{CHO}$  according to known procedures. This reaction is typically performed in the presence of a reducing agent such as sodium cyanoborohydride in a suitable solvent such as methanol.

**[0092]** A compound of formula (I) of this invention wherein X is  $\text{CO}_2$  may be prepared by reacting compound (VII) with either a dicarbonate of formula  $(\text{R}^2\text{OCO})_2\text{O}$  or a chloroformate of formula  $\text{R}^2\text{OCOCl}$  in the presence of a base such as triethylamine in a suitable solvent such as THF at about room temperature for about 1 to 24 hours, preferably under nitrogen.

#### General Reaction Scheme B

**[0093]** A compound of formula (Ib) (i.e., a compound of formula (I) wherein X is a covalent bond and  $\text{R}^2$  is an optionally substituted ring moiety) may be prepared according to the procedures illustrated in Scheme B.



25

[0094] A benzyl acid of formula (VIII) is reacted with an amine of formula (IX) to give an amide, which is then reduced (this step not shown) to give thephenethyl amine of formula (X) according to known procedures (see, for example, Nagarajan et al., *Ind. J. Chem.*, 24B:83-97 (1985)). Compound (X) thus obtained is reacted with a carboxylic acid of formula (XI) to give the benzamide of formula (XII). The compound of formula (XII) is subjected to cyclodehydration followed by reduction, preferably with  $\text{NaBH}_4$  in methanol, to give the tetrahydroisoquinoline (Ib).

30

[0095] The reaction of compound (X) with compound (XI) is carried out in the presence a base such as triethylamine, in a reaction inert solvent such as dichloromethane, THF,  $\text{CHCl}_3$ , or dioxane, preferably in the presence of a coupling agent such as 1-propanephosphonic acid cyclic anhydride (PPAA), EDC, or a mixture thereof, and a catalytic amount of an additive such as HOBt or 4-dimethylaminopyridine (DMAP), at about  $0^\circ\text{C}$  to room temperature for about 1 to 36 hours.

35

[0096] The cyclodehydration of compound (XII) may be carried out according to procedures analogous to those described for in Scheme A.


[0097] Alternatively, compound (X) may be reacted with an acyl chloride of formula (IV) to give an amide (XII). This reaction may be performed according to procedures analogous to those described in Scheme A.

40

[0098] Compound (XI) may be substituted at its  $\text{R}^1$  moiety with a desired substituent. For example, when  $\text{R}^1$  is aryl aldehyde, compound (XI) is coupled with a carboalkoxytriphenylphosphorane in the presence of a base such as sodium hydroxide in a reaction inert solvent such as tetrahydrofuran (THF). This reaction is performed at about room temperature for about 1 to 12 hours.

#### 45 General Reaction Scheme C

[0099] Scheme C illustrates methods to produce a compound of formula (Ic), namely a compound of formula (I) wherein  $\text{A}_1$  is hydroxy and X is a covalent bond,  $(\text{CH}_2)_n$ , or  $(\text{CH}_2)_x$ -phenyl (wherein x is 0 or 1), starting from a compound of formula (Ia). In this scheme, "Pro" is a hydroxy protecting group.



Scheme C

[0100] For example, compound (Ia) wherein  $A_1$  is hydroxy, X is CO, and  $R^2$  is  $CF_3$  is first protected with a suitable hydroxy protecting group to obtain the compound of formula (XIII), which is converted to the compound of formula (XIV), which is coupled with a suitable acyl chloride of formula  $R^2COCl$  to obtain the compound of formula (XV), which is finally deprotected to yield the desired compound (Ic).

[0101] While any hydroxy protecting groups known to those skilled in the art may be used to protect compound (Ia), a benzyl or methyl group is preferably used. The protection is typically carried out by reacting compound (Ia) wherein  $A_1$  is hydroxy with benzyl bromide in the presence of sodium hydride in a reaction inert solvent such as DMF at about  $0^\circ C$  to  $100^\circ C$  under nitrogen for about 1 to 12 hours, preferably about 4 hours.

[0102] Conversion of compound (XIII) is performed according to known procedures. This conversion is conveniently carried out in the presence of an inorganic base such as  $K_2CO_3$ , in a reaction inert solvent such as methanol or ethanol under  $N_2$  for about 1 to 24 hours, preferably about 12 to 24 hours.

[0103] The coupling reaction of compound (XIV) with a desired acyl chloride of formula  $R^2COCl$  is preferably performed in the presence of triethylamine in THF at about room temperature under nitrogen for about 1 to 24 hours.

[0104] The hydroxy protecting group of compound (XV) may be removed by known procedures. For example, this reaction may be performed in the presence of a source of  $H_2$  and a catalyst such as 20%  $Pd(OH)_2/C$ , in a reaction inert solvent such as methanol, under nitrogen, at about the reflux temperature of the reaction mixture for about 1 to 4 hours.

[0105] Compound (I) wherein  $R^1$  is aryl substituted by an alkenylamide group may be obtained by reacting compound (I) wherein  $R^1$  is iodophenyl with a suitable alkenylamide. This reaction may be performed in the presence of a catalyst such as  $Pd(PPh_3)_4$  and a base such as triethylamine in a reaction inert solvent such as N,N-dimethylformamide (DMF) at about  $60^\circ C$  to  $100^\circ C$  for about 1 to 12 hours. The alkenylamide may be prepared by known methods, for example, by reacting a secondary amine with an acyl halide such as acryloyl chloride. This reaction is performed in the presence of a base such as triethylamine in a reaction inert solvent such as dichloroethane at about room temperature for about 1 to 12 hours.

[0106] Compound (I) wherein  $A_1$  is hydroxy may be obtained by converting compound (Ia) wherein  $A_1$  is methoxy. This conversion is performed in the presence of boron tribromide in a suitable solvent such as dichloromethane at about  $-78^\circ C$  to room temperature under nitrogen for about 1 to 12 hours. If required, this conversion may be quenched by adding a suitable solvent such as methanol.


[0107] The hydroxy group at the 6-position of the tetrahydroisoquinoline ring of compound (I) may be converted into an optionally substituted  $(C_1-C_4 \text{ alkyl})-CO$ . For example, the hydroxy may be coupled with 1-(2-chloroethyl)pyrrolidine hydrochloride in the presence of a condensing agent such as sodium hydride in a reaction inert solvent such as DMF. This reaction is preferably performed at about room temperature to the reflux temperature of the reaction mixture, more preferably at about  $100^\circ C$ , under nitrogen, for about 1 to 8 hours, preferably about 4 to 5 hours.

[0108] A substituent attached to a ring moiety of R<sup>1</sup> of compound (I) may be converted to another substituent. For example, the hydroxy substituent on the R<sup>1</sup> ring moiety may be converted to 2-pyrrolidin-1-yl-ethoxy according to procedures analogous to those for the coupling reaction of hydroxy at the 6-position of the tetrahydroisoquinoline ring of compound (I) with 1-(2-chloroethyl)pyrrolidine hydrochloride.

[0109] Compounds (II), (III), (IV), (VIII), (IX) and (XI) are known compounds and are readily prepared according to procedures well known to those skilled in the art.

#### General Reaction Scheme D


[0110] A compound of formula (XXI) (i.e., a compound of formula (II) lacking the R<sup>2</sup> group) may be prepared according to the procedures illustrated in Scheme D.



**Scheme D**

[0111] Compounds of formula (XXI) can be prepared under standard reaction conditions analogous to those well known in the art (e.g., Grethe et al., *J. Org. Chem.*, 33(2):491 (1968)). For example a suitably protected dihydro-1H-isoquinolin-4-one of formula (XXII) can be treated with a nucleophile of formula (XXIII) such as substituted phenyl lithium to give the alcohol of formula (XXIV). The reaction is typically run in an inert solvent such as THF, 1,2 dimethoxyethane, or diethyl ether, and a temperature range of -78°C to room temperature is preferred.


[0112] The alcohol (XXIV) can be reduced to the tetrahydroisoquinoline of formula (XXV) under suitable conditions (e.g., hydrogenation over a metal catalyst such as palladium). Another option for the reaction is to perform the reduction with a trialkylsilane in the presence of a strong acid such as trifluoroacetic acid. The reaction is typically run in an inert solvent at a temperature range of 0°C to 60°C.




[0113] The compounds of formula (XXI) can be treated with an acid chloride of formula (XXVI) or an acid anhydride of formula (XXVII) in the presence a tertiary amine base such as triethylamine or DMAP to give an amide of formula (XXVIII). The acid chlorides of formula (XXVI) and acid anhydride of formula (XXVII) are commercially available or can be prepared from carboxylic acids by procedures known to those skilled in the art.



[0114] Another synthetic route for preparing the compounds of formula (XXVIII) involves treating a compound of formula (XXI), preferably at room temperature, with a carboxylic acid of formula (XXIX) in the presence of a coupling agent (e.g., dicyclohexylcarbodiimide (DCC), 1-(3'-dimethylaminopropyl)-3-ethylcarbodiimide (EDC), or 1-propanephosphonic acid cyclic anhydride (PPAA)) and a suitable base (e.g., triethylamine, DMAP, or N-methylmorpholine (NMO)) in a solvent such as dichloromethane, chloroform, or dimethylformamide. Optionally, agents such as HOBT may be added to the reaction.




[0115] Compounds of formula (XXX) can be prepared by reacting the amines of formula (XXI) with a sulfonyl chloride of formula (XXXI) in the presence of a base such as triethylamine, DMAP, or NMO in a suitable solvent such as dichloromethane, chloroform, or dimethylformamide. The reactions can be run in temperature range of -20°C to 60°C, though room temperature is preferred.



[0116] Compounds of formula (XXXII) can be prepared by reacting the amines of formula (XXI) with a chloroformate of formula (XXXIII) in the presence of a base such as triethylamine, DMAP, NMO, or sodium hydrogen carbonate in a suitable solvent such as dichloromethane, chloroform, aqueous or anhydrous tetrahydrofuran, or dimethylformamide. The reaction can be run at 0°C to 60°C, though room temperature is preferred.

#### General Reaction Scheme E

[0117] A compound of formula (XXI) (i.e., a compound of formula (II) lacking the R² group) may also be prepared according to the procedures illustrated in Scheme E.




Scheme E

[0118] An alternative and more preferable synthesis of compounds of formula (XXI) involves reacting an alpha halogenated ketone of formula (XXXIV) with a suitably protected ammonia equivalent, preferably benzylamine, to give an amino-ketone of formula (XXXV). It will be obvious to anyone skilled in the art that the order of some of the next steps may be reversed. The compound of formula (XXXV) can be reduced with a suitable reducing agent, preferably sodium borohydride, di-isobutyl aluminum hydride (DIBAL-H), or lithium aluminum hydride in a suitable solvent. Typically, sodium borohydride reactions are run in protic solvents such as methanol or ethanol, in the temperature range of  $-10^\circ\text{C}$  to  $40^\circ\text{C}$  (but preferably at about  $0^\circ\text{C}$ ), and the DIBAL-H or lithium aluminum hydride reactions are preferably run in dichloromethane or THF in the temperature range of  $-78^\circ\text{C}$  to  $30^\circ\text{C}$  (preferably at about  $0^\circ\text{C}$ ) to give the amino alcohol of formula (XXXVI). In the next step, the amino alcohol of formula (XXXVI) is condensed with an aldehyde of formula (XXXVII) to give initially an iminium salt of formula (XXXVIII). The iminium salt of formula (XXXVIII) is reduced in situ with a reducing agent such as sodium borohydride, sodium cyanoborohydride, or sodium triacetoxyborohydride in a solvent such as dichloromethane, chloroform, 1,2 dichloroethane, methanol, or ethanol to give the amino alcohol of formula (XXXIX). For the latter reduction reaction, a temperature range of  $-10^\circ\text{C}$  to  $50^\circ\text{C}$  can be employed but a temperature of about  $0^\circ\text{C}$  is preferred.

[0119] The amino alcohol of formula (XXXIX) can be cyclodehydrated in the presence of a strong acid such as sulfuric acid or trifluoroacetic acid in a solvent such as dichloromethane, chloroform, or 1,2 dichloroethane in a temperature range of  $0^\circ\text{C}$  to  $60^\circ\text{C}$  to give (after appropriate deprotection) the tetrahydroisoquinoline of formula (XXI).

#### General Reaction Scheme F

[0120] A compound of formula (XXIX) (i.e., the amino alcohol intermediary shown in Scheme E) may be prepared according to the alternative procedures illustrated in Scheme F.



Scheme F

[0121] Another option for the synthesis of amino alcohol of formula (XXXIX) is via the amide of formula (XL), which can be prepared by methods known to those of ordinary skill from the amino alcohol (XXXVI). Such methods may include taking the amino-alcohol of formula (XXXVI) and treating this with an acid chloride of formula (XLI) in the presence of a tertiary amine base such as triethylamine or DMAP. The acid chlorides of formula (XLI) are commercially available or can be prepared from carboxylic acids by procedures known to those of ordinary skill in the art. Another method of preparing the amides of formula (XL) involves treating an amino alcohol of formula (XXXVI) with a carboxylic acid of formula (XLII) in the presence of a coupling agent (e.g., DCC, EDC, or PPAA) and a suitable base (e.g., triethylamine, DMAP, or NMO) in a solvent such as dichloromethane, chloroform, or dimethylformamide. Optionally, agents such as HOBt may be added to the reaction. Typically, the reaction is run in the temperature range of 0°C to 50°C, with room temperature being preferred.

[0122] The amides of formula (XL) can then be reduced to an amino-alcohol of formula (XXXIX) by reaction with a reducing agent such as lithium aluminum hydride or borane in a suitable solvent such as tetrahydrofuran or 1,2 dimethoxyethane at -10°C to 100°C (with 0°C being preferred).

[0123] Reactive groups not involved in the above processes can be protected with standard protecting groups during the reactions and removed by standard procedures (Greene & Wuts, Protecting Groups in Organic Synthesis, John Wiley and Sons, Inc., Interscience, 2nd edition, New York) known to those of ordinary skill in the art. Presently preferred protecting groups include methyl and benzyl for the hydroxyl moiety, and trifluoroacetamide and benzyl for the amino moiety.

[0124] In each of the general reaction schemes discussed or illustrated above, pressure is not critical unless otherwise indicated. Pressures from about 0.5 atmospheres (~50,000 Pa) to about 5 atmospheres (~500,000 Pa) are generally acceptable, and ambient pressure, i.e. about 1 atmosphere (~100,000 Pa), is preferred as a matter of convenience.

[0125] The compounds of this invention may be prepared in racemic form and be resolved into their component enantiomers by standard techniques such as fractional crystallization or preparative chromatography. Alternatively, enantiomers of the invention compounds may be synthesized from the appropriate optically active intermediates or starting materials using any of the general processes described herein.

[0126] Further, optically active compounds of this invention may be prepared using enantioselective reactions or by resolution techniques such as the preparation of diastereomers by reacting the racemic material with an optically active reagent.

[0127] The compounds of this invention are valuable estrogen agonists or antagonists, and thus valuable pharmaceutical agents. Those that are estrogen agonists are useful for oral contraception, relief of the symptoms of menopause, prevention of threatened or habitual abortion, relief of dysmenorrhea, relief of dysfunctional uterine bleeding, relief of endometriosis, an aid in ovarian development, treatment of acne, diminution of excessive growth of body hair in women (hirsutism), the prevention and/or treatment of cardiovascular disease, prevention and treatment of athero-

sclerosis, prevention and treatment of osteoporosis, treatment of benign prostatic hyperplasia and prostatic carcinoma, obesity, and suppression of post-partum lactation. These agents also have a beneficial effect on plasma lipid levels and as such are useful in treating and/or preventing hypercholesterolemia.

[0128] While the compounds of this invention are typically estrogen agonists in bone, unexpectedly they are also often antiestrogens in breast tissue and as such would be useful in the treatment and prevention of breast cancer.

[0129] While the compounds of this invention are expected to be sufficiently active if administered separately, in many instances it will be beneficial to combine these compounds with other compounds in order to even more effectively treat a disease or condition. Exemplary categories of compounds which will be beneficial in combination with the compounds of the present invention include an anabolic agent, a prodrug thereof, or a pharmaceutically acceptable salt of said anabolic agent or said prodrug; a growth hormone or a growth hormone secretagogue, a prodrug thereof, or a pharmaceutically acceptable salt of said growth hormone secretagogue or said prodrug; a prostaglandin agonist/antagonist, a prodrug thereof, or a pharmaceutically acceptable salt of said prostaglandin agonist/antagonist or said prodrug; or a parathyroid hormone or sodium fluoride.

[0130] It will also be useful to present the compounds of the present invention in the form of kits useful for treating or preventing a disease, disorder, condition, or symptom mediated by an estrogen receptor and/or caused by lowered estrogen levels, said kit comprising a compound of the present invention and a pharmaceutically acceptable vehicle, carrier, or diluent in a dosage form, and a container for containing said dosage form.

[0131] In many instances, it will be preferable for the kit to include, in addition to a compound of the present invention, an anabolic agent, a prodrug thereof, or a pharmaceutically acceptable salt of said anabolic agent or said prodrug; a growth hormone or a growth hormone secretagogue, a prodrug thereof, or a pharmaceutically acceptable salt of said growth hormone secretagogue or said prodrug; a prostaglandin agonist/antagonist, a prodrug thereof, or a pharmaceutically acceptable salt of said prostaglandin agonist/antagonist or said prodrug; or a parathyroid hormone or sodium fluoride.

#### Methods for Testing the Activity of Invention Compounds

##### Assay 1: Control and Prevention of Endometriosis

[0132] A preferred protocol for surgically inducing endometriosis is that described by Jones, *Acta Endocrinol (Copenh)*, 106:282-88 (1984). Adult Charles River Sprague-Dawley CD® female rats (200-240 g) are used. An oblique ventral incision is made through the skin and musculature of the body wall. A segment of the right uterine horn is excised, the myometrium is separated from the endometrium, and the segment is cut longitudinally. A 5x5 mm section of the endometrium, with the epithelial lining apposed to the body wall, is sutured at its four corners to the muscle using polyester braid (Ethiflex, 7-0®). The criterion of a viable graft is the accumulation of fluid similar to that which occurs in the uterus as a result of estrogen stimulation.

[0133] Three weeks after transplantation of the endometrial tissue (+3 weeks) the animals are laparotomized, the volume of the explant (length x width x height) in mm is measured with calipers, and treatment is begun. The animals are injected sc for 3 weeks with 10 to 1000 mg/kg/day of a test compound according to the present invention. Control animals bearing endometrial explants are injected sc with 0.1 ml/day of corn oil for 3 weeks. At the end of the 3 week treatment period (+6 weeks), the animals are laparotomized and the volume of the explant determined. Eight weeks after cessation of treatment (+14 weeks) the animals are sacrificed and the explants are measured again. Statistical analysis of the explant volume is by an analysis of variance.

##### Assay 2: Effect on Prostate Weight

[0134] Three-month-old male Sprague-Dawley rats are administered by subcutaneous injection control vehicle (10% ethanol in water), estradiol (30 µg/kg), testosterone (1 mg/kg), or a test compound according to the present invention daily for 14 days (n=6/group). After 14 days the animals are sacrificed, the prostate is removed, and the wet prostate weight is determined. Mean weight is determined and statistical significance (p<0.05) is determined compared to the vehicle-treated group using the Student's t-test.

[0135] Active compounds significantly (P<0.05) decrease prostate weight compared to controls. Testosterone is expected to have no effect while estrogen at 30 µg/kg is expected to significantly reduce prostate weight.

##### Assay 3: In Vitro Estrogen Receptor Binding

[0136] An *in vitro* estrogen receptor binding assay that measures the ability of the compounds of the present invention to displace [<sup>3</sup>H]-estradiol from human estrogen receptor obtained by recombinant methods in yeast, bacteria, or mammalian cells is used to determine the estrogen receptor binding affinity of the compounds of this invention. The materials

used in this assay are: (1) TD-0.3 assay buffer (containing 10 mM Tris, pH 7.6, 0.3 M potassium chloride, and 5mM DTT); (2) the radioligand used is [<sup>3</sup>H]-estradiol obtained from New England Nuclear (Boston, MA); (3) the cold ligand used is estradiol obtained from Sigma (St. Louis, MO); and (4) recombinant human estrogen receptor, hER.

**[0137]** A solution of the compound being tested is prepared in TD-0.3 with 4% DMSO and 16% ethanol. The tritiated estradiol is dissolved in TD-0.3 such that the final concentration in the assay is 5 nM. The hER is also diluted with TD-0.3 such that 0.2 nM hER is in each assay well. Using microtitre plates, each incubate receives 50 µl of cold estradiol (nonspecific binding) or the test compound solution, 20 µl of the tritiated estradiol, and 30 µl of hER solutions. Each plate contains varying concentrations of the compound and total binding controls in triplicate. The plates are incubated overnight at 4°C. The binding reaction is then terminated by the addition and mixing of 100 µl of 3% hydroxylapatite in 10 mM Tris, pH 7.6, followed by incubation for 15 minutes at 4°C. The mixture is centrifuged and the pellet washed four times with 1% Triton-X100 in 10 mM Tris, pH 7.6. The hydroxylapatite pellets are suspended in Ecolite (+) (ICN Biomedicals, Inc., Aurora, OH) and radioactivity is assessed using beta scintigraphy. The mean of all triplicate data points (counts per minute, cpm's) is determined. Specific binding is calculated by subtracting nonspecific cpm's (defined as counts that remain following separation of reaction mixture containing recombinant receptor, radioligand, and excess unlabeled ligand) from total bound cpm's (defined as counts that remain following the separation of reaction mixture containing only recombinant receptor, radioligand). Test compound potency is determined by means of IC<sub>50</sub> determinations (the concentration of a test compound needed to inhibit 50% of the of the total specific tritiated estradiol bound). Specific binding in the presence of varying concentrations of test compound is determined and calculated as percent specific binding of total specific radioligand bound. Data are plotted as percent inhibition by test compound (linear scale) versus test compound concentration (log scale).

#### Assay 4: Effect on Total Cholesterol Levels

**[0138]** The effect of the compounds of the present invention on plasma levels of total cholesterol is measured as follows. Blood samples are collected via cardiac puncture from anesthetized female (Sprague-Dawley) rats 4-6 months of age that are bilaterally ovariectomized and treated with the test compound (10-1000 µg/kg/day, for example, sc or orally for 28 days or with control vehicle for the same time), or sham operated. The blood is placed in a tube containing 30 µl of 5% EDTA (10 µl EDTA/1 ml of blood). After centrifugation at 2500 rpm for 10 minutes at 20°C the plasma is removed and stored at -20°C. The total cholesterol is assayed using a standard enzymatic determination kit from Sigma Diagnostics (Procedure No. 352).

#### Assay 5: Effect on Obesity

**[0139]** Ten-month-old female Sprague-Dawley rats, weighing approximately 450 grams, are sham-operated or ovariectomized and treated orally with vehicle, 17α-ethynyl-estradiol at 30 mg/kg/day, or a compound according to the present invention at 10-1000 mg/kg/day for 8 weeks. There are 6 to 7 rats in each sub group. On the last day of the study, body composition of all rats is determined via dual energy x-ray absorptiometry using a Hologic QDR-1000/W (Hologic, Bedford, MA) equipped with whole body scan software that shows the proportions of fat body mass and lean body mass.

**[0140]** A decrease in fat body mass indicates that the test compound is useful in preventing and treating obesity.

#### ABBREVIATIONS

**[0141]** Abbreviations used in the following examples and preparations include:

1,2 DCE	1,2-Dichloroethane
d	Doublet
dd	Double Doublet
DMAP	4-Dimethylamino Pyridine
EDC	1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide Hydrochloride
EtOAc	Ethyl Acetate
EtOH	Ethyl Alcohol or Ethanol
Et <sub>2</sub> O	Ethyl Ether
Et <sub>3</sub> N	Triethylamine
HOBt	1-Hydroxybenzotriazole
HPLC	High Pressure Liquid Chromatography
hr	Hour(s)
m	Multiplet

MeOH	Methyl Alcohol or Methanol
min	Minute(s)
MS	Mass Spectrometry
NMR	Nuclear Magnetic Resonance
5 PPAA	1-Propanephosphonic Acid Cyclic Anhydride
q	Quartet
RT (or rt)	room temperature (about 20-25° C)
s	Singlet
sat.	Saturated
10 t	Triplet
TBAF	Tetrabutyl Ammonium Fluoride
TFA	Trifluoroacetic Acid
THF	Tetrahydrofuran

## 15 EXAMPLES AND PREPARATIONS

[0142] The following examples will serve to illustrate, but do not limit, the invention that is defined by the claims.

### Preparation 1

#### 20 4-Benzyloxy-N-[2-(3-methoxyphenyl)ethyl]benzamide

[0143] A mixture of 3-methoxyphenethylamine (16.557 g, 109.5 mmol), 4-benzyloxybenzoic acid (25.000 g, 109.5 mmol), 1-hydroxybenzotriazole hydrate (22.202 g, 164.3 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (22.047 g, 115.0 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (300 ml) was stirred at 0°C under N<sub>2</sub> for 1 hr, then warmed to rt and stirred at rt for 19 hr. The reaction mixture was washed sequentially with 1M NaOH (150 ml), 1M HCl (150 ml), and H<sub>2</sub>O (150 ml), then dried over MgSO<sub>4</sub> and concentrated *in vacuo* to give 39.00 g (99% yield) of yellow solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.63 (d, J = 8.7 Hz, 2H), 7.43 - 7.30 (m, 5H), 7.21 (d, J = 7.7 Hz, 1H), 6.95 (d, J = 8.7 Hz, 2H), 6.82 - 6.78 (m, 2H), 6.76 (s, 1H), 6.02 (m, 1H), 5.08 (s, 2H), 3.77 (s, 3H), 3.68 (dd, J = 13.07 Hz, J = 6.64 Hz, 2H), 2.90 - 2.86 (m, 2H); MS m/e 362 (M<sup>+</sup>+1).

### Preparation 2

#### 35 1-(4-Benzyloxyphenyl)-6-methoxy-3,4-dihydroisoquinoline hydrochloride

[0144] Phosphorus pentachloride (58.557 g, 281.2 mmol) was added in portions to a solution of 4-benzyloxy-N-[2-(3-methoxyphenyl)ethyl]benzamide (59.775 g, 165.4 mmol) in 1,2-dichloroethane (500 ml) at 0°C under N<sub>2</sub>. The resulting reaction was stirred at 0°C for 30 min, then refluxed for 18 hr. After cooling tort, hexane (1200 ml) was added and the resulting suspension was cooled to 0°C. The solvent was decanted off from the oil that separated and the remaining oil was dried *in vacuo* to give 54.56 g (87% yield) of orange residue. MS m/e 344 (M<sup>+</sup>+1 - HCl).

### Preparation 3

#### 45 1-(4-Benzyloxyphenyl)-6-methoxy-1,2,3,4-tetrahydroisoquinoline

[0145] Sodium borohydride (10.874 g, 287.3 mmol) was added in small portions to a solution of 1-(4-benzyloxyphenyl)-6-methoxy-3,4-dihydroisoquinoline (54.56 g, 143.6 mmol) in MeOH (600 ml) at 0°C. The reaction mixture was stirred at 0°C for 1 hr, then quenched by the dropwise addition of H<sub>2</sub>O (20 ml). The resulting mixture was concentrated *in vacuo* to a volume of about 100 ml. The solid that separated was collected to give 35.926 g (72% yield) of white solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.42 - 7.28 (m, 5H), 7.16 (d, J = 8.72 Hz, 2H), 6.91 (d, J = 8.72 Hz, 2H), 6.67 - 6.58 (m, 2H), 6.64 (s, 1H), 5.03 (s, 2H), 4.98 (s, 1H), 3.76 (s, 3H), 3.26 - 3.21 (m, 1H), 3.08 - 2.95 (m, 2H), 2.79 - 2.74 (m, 1H); MS m/e 346 (M+1).

[0146] The remaining filtrate was evaporated *in vacuo* to give an additional 23.85 g of crude material.

**Preparation 4****1-[1-(4-Benzyloxyphenyl)-6-methoxy-3,4-dihydro-1H-isoquinolin-2-yl]-2,2,2-trifluoroethanone**

[0147] Trifluoroacetic anhydride (22.620 g, 107.7 mmol) was added to a solution of 1-(4-benzyloxyphenyl)-6-methoxy-1,2,3,4-tetrahydroisoquinoline (28.530 g, 82.6 mmol) and Et<sub>3</sub>N (15.0 ml, 107.7 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (500 ml) at 0°C under N<sub>2</sub>. The resulting yellow solution was stirred at 0°C for 3 hr, diluted with CH<sub>2</sub>Cl<sub>2</sub> (150 ml), and washed first with 1 M HCl (2 x 100 ml) then 1 M NaOH (100 ml), dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to give 32.09 g of yellow oil. Purification by flash chromatography, eluting with hexane : EtOAc (9:1) gave 26.97 g (74% yield) of yellow oil.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.40 - 7.29 (m, 5H), 7.11 (d, J = 8.72 Hz, 2H), 6.94 (d, J = 8.51 Hz, 1H), 6.87 (d, J = 8.72 Hz, 2H), 6.77 - 6.69 (m, 3H), 5.01 (s, 2H), 3.91 - 3.80 (m, 1H), 3.79 (s, 3H), 3.48 - 3.41 (m, 1H), 3.10 - 3.02 (m, 1H), 2.85 - 2.80 (m, 1H); MS m/e 442 (M<sup>+</sup>+1).

**Preparation 5****1-{6-Benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinolin-2-yl}-2,2,2-trifluoroethanone**

[0148] A solution of 2,2,2-trifluoro-1-{6-hydroxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinolin-2-yl}ethanone (1.886 g, 4.3 mmol) in anhydrous DMF (50 ml) was added to a suspension of NaH (0.104 g, 4.3 mmol) in anhydrous DMF (100 ml) at rt under N<sub>2</sub>. After stirring at rt for 1 hr, a solution of benzyl bromide (0.782 g, 4.6 mmol) in anhydrous DMF (10 ml) was added and the reaction mixture was heated to 100°C for 4 hr. The reaction mixture was cooled, diluted with H<sub>2</sub>O (250 ml), and extracted with EtOAc (4 x 100 ml). The combined extracts were washed with H<sub>2</sub>O (2 x 100 ml), dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to give 2.71 g of orange oil.

Purification by flash chromatography, eluting with EtOAc : MeOH (7:3) gave 0.745 g (33% yield) of yellow oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.42 - 7.39 (m, 4H), 7.38 - 7.33 (m, 1H), 7.20 - 7.10 (m, 2H), 6.95 - 6.92 (m, 1H), 6.85 - 6.79 (m, 4H), 6.72 (or s, 1H), 5.05 (s, 2H), 4.30 - 4.18 (m, 2H), 3.92 - 3.88 (m, 1H), 3.47 - 3.40 (m, 1H), 3.30 - 2.70 (m, 8H), 1.94 (br s, 4H); MS m/e 525 (M<sup>+</sup>+1).

**Preparation 6****6-Benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinoline**

[0149] A mixture of 1-{6-benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinolin-2-yl}-2,2,2-trifluoroethanone (1.088 g, 2.1 mmol) and anhydrous K<sub>2</sub>CO<sub>3</sub> (2.861 g, 20.7 mmol) in MeOH (150 ml) was refluxed under N<sub>2</sub> for 18 hr, then evaporated *in vacuo* to a yellow residue. This was dissolved in H<sub>2</sub>O (25 ml) and extracted with EtOAc (3 x 50 ml). The combined extracts were dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to give 0.881 g of crude yellow product. Purification by flash chromatography, eluting first with EtOAc : MeOH (8:2), then with EtOAc : MeOH (1:1) gave 0.842 g (95% yield) of yellow oil.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.41 - 7.29 (m, 5H), 7.17 - 7.12 (m, 2H), 6.86 - 6.52 (m, 5H), 5.07 - 5.05 (m, 1H), 5.02 - 4.98 (m, 2H), 4.12 - 4.10 (m, 2H), 3.24 - 3.21 (m, 1H), 3.06 - 2.95 (m, 1H), 2.91 - 2.90 (m, 3H), 2.78 - 2.75 (m, 1H), 2.65 (br s, 4H), 1.81-1.80 (m, 4H); MS m/e 429 (M<sup>+</sup>+1).

**Preparation 7****{6-Benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-3,4-dihydro-1H-isoquinolin-2-yl}phenylmethanone**

[0150] Benzoyl chloride (0.013 g, 0.09 mmol) was added to a solution of 6-benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinoline (0.040 g, 0.09 mmol) and Et<sub>3</sub>N (0.026 ml, 0.19 mmol) in anhydrous THF (10 ml) at rt under N<sub>2</sub>. The resulting suspension was stirred at rt for 20 hr, then evaporated *in vacuo* to give 0.091 g of white solid. Purification by flash chromatography, eluting with EtOAc : MeOH (6:4) gave 0.049 g (98% yield) of white solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.44 - 7.32 (m, 10H), 7.00 - 6.96 (m, 2H), 6.82 - 6.79 (m, 6H), 5.05 (m, 2H), 4.22 - 4.20 (m, 2H), 3.62 - 3.58 (m, 1H), 3.32 - 3.26 (m, 2H), 3.11-3.09 (m, 2H), 2.96 - 2.92 (m, 4H), 2.69 - 2.65 (m, 1H), 1.92 br s, 4H; MS m/e 533 (M<sup>+</sup>+1).

**Preparation 8****1-[6-Benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinolin-2-yl]-2,2-dimethylpropan-1-one**

**[0151]** A solution of 6-benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinoline (0.015 g, 0.04 mmol) and Et<sub>3</sub>N (0.01 ml, 0.07 mmol) in anhydrous THF (0.35 ml) was added to a solution of trimethylacetyl chloride (0.04 mmol) in anhydrous THF (0.4 ml) at rt in a sealed reaction vessel. The resulting suspension was stirred at rt for 20 hr, then evaporated to dryness. The product was suspended in a mixture of H<sub>2</sub>O (0.4 ml) and saturated NaHCO<sub>3</sub> solution (0.4 ml), then extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 0.75 ml). The combined extracts were evaporated to dryness to give the crude product. Purification by reverse-phase HPLC, eluting first with gradient. Purification by reverse-phase HPLC on a Primesphere (Phenomenex 2320 West 205<sup>th</sup> St. Torrance, CA 90501) C-18HC (50.0 mm x 10.0 mm column with 5 μm particle size) column, eluting with a linear gradient starting at time 0 min. of H<sub>2</sub>O : CH<sub>3</sub>CN : 1%TFA/H<sub>2</sub>O (85 : 10 : 5), increasing to H<sub>2</sub>O : CH<sub>3</sub>CN : 1%TFA/H<sub>2</sub>O (5 : 90 : 5) at 8 min., detected on a Micromass Platform 2 mass spectrometer (DAD 190 - 600 nM) gave material which, after evaporation to dryness, was purified by reverse-phase HPLC on a Primesphere C-18HC (3.0 mm x 2.0 mm column with 5 μm particle size) column, eluting with a linear gradient starting at time 0 min. of H<sub>2</sub>O : CH<sub>3</sub>CN : TFA (99.9 : 0 : 0.1), increasing to H<sub>2</sub>O : CH<sub>3</sub>CN : TFA (0 : 99.9 : 0.1) at 4 min., detected with a UV detector (300 nM +/- 90 nM and 254 nM +/- 25 nM), gave an eluent that was evaporated to dryness to give 0.012 g (56% yield) of the pure product. MS 513 m/e (M<sup>+</sup>+1).

**Preparation 9****4-Benzyloxy-N-phenethylbenzamide**

**[0152]** A mixture of phenethylamine (5.000 g, 41.3 mmol), 4-benzyloxybenzoic acid (9.427 g, 41.3 mmol), 1-hydroxy-benzotriazole hydrate (8.378 g, 62.0 mmol) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (8.320 g, 43.4 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (100 ml) was stirred at 0°C under N<sub>2</sub> for 1 hr, then warmed to rt and stirred at rt for 22 hr. The reaction mixture was washed sequentially with 1 M NaOH (50 ml), 1 M HCl (50 ml), and H<sub>2</sub>O (50 ml), then dried over MgSO<sub>4</sub> and concentrated *in vacuo* to give 13.61 g of tan solid. Purification by flash chromatography, eluting first with hexane : EtOAc (1:1) and then with EtOAc gave 12.20 g (89% yield) of white solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.63 (d, J = 8.92 Hz, 2H), 7.42 - 7.36 (m, 4H), 7.35 - 7.29 (m, 3H), 7.24 - 7.21 (m, 3H), 6.95 (d, J = 8.92 Hz, 2H), 6.00 (br s, 1H), 5.08 (s, 2H), 3.71 - 3.66 (m, 2H), 2.92 - 2.89 (m, 2H); MS m/e 332 (M<sup>+</sup>+1).

**Preparation 10****1-(4-Benzyloxyphenyl)-3,4-dihydroisoquinoline**

**[0153]** A solution of 4-benzyloxy-N-phenethylbenzamide (1.000 g, 3.0 mmol) in POCl<sub>3</sub> (10 ml, 107.3 mmol) was refluxed under N<sub>2</sub> for 18 hr, then the cooled reaction mixture was slowly added to ice-water (50 ml) with stirring. The resulting mixture was extracted with EtOAc (4 x 20 ml), then the remaining aqueous layer was basified to pH 10 with concentrated ammonia (80 ml). This was extracted with EtOAc (3 x 50 ml), then the combined extracts were dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to give 0.270 g of yellow solid. Purification by flash chromatography, eluting with hexane : EtOAc (6:4) gave 0.151 g (16% yield) of yellow solid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.55 (d, J = 8.92 Hz, 2H), 7.45 - 7.43 (d, J = 7.06 Hz, 2H), 7.40 - 7.29 (m, 6H), 7.26 - 7.24 (m, 1H), 7.00 (d, J = 8.92 Hz, 2H), 5.11 (s, 2H), 3.81 - 3.78 (m, 2H), 2.79 - 2.75 (m, 2H) MS m/e 314 (M<sup>+</sup>+1).

**Preparation 11****1-(4-Benzyloxyphenyl)-1,2,3,4-tetrahydroisoquinoline**

**[0154]** Sodium borohydride (0.34 g, 0.89 mmol) was added in small portions to a solution of 1-(4-benzyloxyphenyl)-3,4-dihydroisoquinoline (0.139 g, 0.44 mmol) in MeOH (10 ml) at 0°C. The reaction mixture was stirred at 0°C for 30 min, stirred at rt for 1 hr, then quenched by the dropwise addition of H<sub>2</sub>O (1.0 ml). The resulting mixture was concentrated *in vacuo* to remove the methanol, and the remaining aqueous layer was extracted with CHCl<sub>3</sub> (4 x 10 ml). The combined extracts were dried over MgSO<sub>4</sub> and concentrated *in vacuo* to give 0.138 g (99% yield) of colorless oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.43 - 7.29 (m, 5H), 7.18 - 7.16 (m, 2H), 7.13 - 7.12 (m, 2H), 7.10 - 7.00 (m, 1H), 6.92 (d, J = 8.72 Hz, 2H), 6.75 (d, J = 7.68, 1H), 5.05 (s, 1H), 5.04 (s, 2H), 3.28 - 3.23 (m, 1H), 3.11 - 2.98 (m, 2H), 2.82 - 2.77 (m, 1H); MS m/e 316 (M<sup>+</sup>+1).



## Preparation 12

## 1-[1-(4-Benzyloxyphenyl)-3,4-dihydro-1 H-isoquinolin-2-yl]-2,2,2-trifluoroethanone

[0155] Trifluoroacetic anhydride (0.088 g, 0.42 mmol) was added to a solution of 1-(4-benzyloxyphenyl)-1,2,3,4-tetrahydroisoquinoline (0.132 g, 0.42 mmol) and Et<sub>3</sub>N (0.117 ml, 0.84 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (5 ml) at 0°C under N<sub>2</sub>. The resulting yellow solution was stirred at rt for 5 hr, washed first with 1M HCl (5 ml) then 1M NaOH (5 ml), dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to give 0.186 g of yellow oil.

Purification by flash chromatography, eluting with hexane : EtOAc (9:1) gave 0.147 g (85% yield) of yellow oil.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.41 - 7.26 (m, 6H), 7.23 - 7.17 (m, 2H), 7.12 (d, J = 8.72 Hz, 2H), 7.03 (d, J = 7.26 Hz, 1H), 6.88 (d, J = 8.72 Hz, 2H), 6.78 (s, 1H), 5.02 (s, 2H), 3.95 - 3.91 (m, 1H), 3.52 - 3.44 (m, 1H), 3.14 - 3.05 (m, 1H), 2.90 - 2.86 (m, 1H); MS m/e 410 (M<sup>+</sup>-1).

## Preparation 13

## Toluene-4-sulfonic acid 4-(6-methoxy-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-1-yl)phenyl ester

[0156] A solution of 2,2,2-trifluoro-1-[1-(4-hydroxyphenyl)-6-methoxy-3,4-dihydro-1H-isoquinolin-2-yl]ethanone (3.387 g, 9.64 mmol), p-toluenesulfonyl chloride (2.206 g, 11.6 mmol) and Et<sub>3</sub>N (1.6 ml, 11.6 mmol) in acetone (50 ml) was refluxed under N<sub>2</sub> for 5 hr, then concentrated *in vacuo*. The remaining residue was suspended in H<sub>2</sub>O (100 ml) and extracted with CHCl<sub>3</sub> (4 x 75 ml). The combined extracts were dried over MgSO<sub>4</sub> and concentrated *in vacuo* to give 5.622 g of orange oil. Purification by flash chromatography, eluting with hexane : EtOAc (8:2) gave 4.383 g (90% yield) of white solid.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.71 - 7.66 (m, 2H), 7.31 - 7.25 (m, 2H), 7.12 (d, J = 8.72 Hz, 1H), 7.12 - 7.03 (m, 1H), 6.98 - 6.88 (m, 3H), 6.78 - 6.68 (m, 2H), 6.70 (s, 1H), 3.91 - 3.87 (m, 1H), 3.80 (s, 3H), 3.49 - 3.43 (m, 1H), 2.95 - 2.91 (m, 1H), 2.84 - 2.80 (m, 1H), 2.43 (d, J = 9.13 Hz, 3H); MS m/e 506 (M<sup>+</sup>+1).

## Preparation 14

## Toluene-4-sulfonic acid 4-(6-hydroxy-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-1-yl)phenyl ester

[0157] A 1M solution of BBr<sub>3</sub> in CH<sub>2</sub>Cl<sub>2</sub> (16.2 ml, 16.2 mmol) was added slowly to a solution of toluene-4-sulfonic acid 4-(6-methoxy-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-1-yl)phenyl ester (7.459 g, 14.8 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (500 ml) at 0°C under N<sub>2</sub>. After stirring at 0°C for 1 hr, the reaction mixture was warmed to rt and stirred at rt for 18 hr. MeOH (250 ml) was slowly added with stirring and the resulting solution was concentrated *in vacuo* to give 9.793 g of brown residue.

Purification by flash chromatography, eluting with hexane : EtOAc (7:3) gave 2.764 g (38% yield) of white solid.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.72 - 7.69 (m, 2H), 7.30 (d, J = 8.51 Hz, 2H), 7.12 (d, J = 8.92 Hz, 2H), 6.95 - 6.85 (m, 3H), 6.72 - 6.67 (m, 3H), 3.93 - 3.89 (m, 1H), 3.40-3.32 (m, 1H), 3.07 - 2.94 (m, 1H), 2.81 - 2.77 (m, 1H), 2.44 (s, 3H); MS m/e 492 (M<sup>+</sup>+1).

## Preparation 15

## Toluene-4-sulfonic acid 4-[6-(2-pyrrolidin-1-yl-ethoxy)-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-1-yl]phenyl ester

[0158] A solution of toluene-4-sulfonic acid 4-(6-hydroxy-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-1-yl)phenyl ester (1.790 g, 3.64 mmol) in anhydrous DMF (30 ml) was added to a suspension of NaH (0.175 g, 7.28 mmol) in anhydrous DMF (30 ml) at rt under N<sub>2</sub>. After stirring at rt for 1 hr, a solution of 1-(2-chloroethyl)pyrrolidine hydrochloride (0.619 g, 3.64 mmol) in anhydrous DMF (20 ml) was added and the reaction mixture was heated to 100°C for 5 hr. The reaction mixture was cooled to rt, diluted with H<sub>2</sub>O (200 ml), and extracted with EtOAc (5 x 50 ml). The combined extracts were washed with H<sub>2</sub>O (2 x 50 ml), dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to give 2.193 g of brown oil. Purification by flash chromatography, eluting first with EtOAc : MeOH (9:1), and then with EtOAc : MeOH (7:3), gave 0.482 g (22% yield) of yellow oil.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.70 (d, J = 8.30 Hz, 2H), 7.31 (d, J = 8.09 Hz, 2H), 7.11 (d, J = 8.51 Hz, 2H), 6.92 - 6.89 (m, 3H), 6.76 - 6.71 (m, 3H), 4.56 - 4.51 (m, 2H), 3.93 - 3.89 (m, 3H), 3.47 (br s, 2H), 3.39 - 3.33 (m, 1H), 3.08 - 2.97 (m, 3H), 2.84 - 2.80 (m, 1H), 2.44 (s, 3H) 2.24 - 2.21 (m, 2H), 2.16 - 2.10 (m, 2H); MS m/e 589 (M<sup>+</sup>+1).

**Preparation 16****4-Iodo-N-[2-(3-methoxyphenyl)-ethyl]benzamide**

5 **[0159]** A mixture of 3-methoxyphenethylamine (11.7 g, 77.4 mmol), 4-iodobenzoyl chloride (20.6 g, 77.4 mmol), and Et<sub>3</sub>N (8.2 g, 81.2 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (250 ml) was stirred at 0°C under N<sub>2</sub> for 1 hr, then warmed to rt and stirred at rt for 19 hr.

The reaction mixture was washed sequentially with 1M HCl (2 x 150 ml), 1M NaOH (2 x 150 ml), and saturated NaCl solution (150 ml), then dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo* to give 21 g (72% yield) of white solid.

10 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.74 (d, J = 8.52 Hz, 2H), 7.39 (d, J = 8.52 Hz, 2H), 7.25 - 7.21 (m, 1H), 6.81 - 6.75 (m, 3H), 6.06 (br s, 1H), 3.78 (s, 3H), 3.71 - 3.67 (m, 2H), 2.91 - 2.87 (m, 2H); MS m/e 382 (M<sup>+</sup>+1).

**Preparation 17****1-(4-Iodophenyl)-6-methoxy-3,4-dihydroisoquinoline hydrochloride**

**[0160]** Phosphorus pentachloride (11 g, 52.8 mmol) was added in portions to a solution of 4-iodo-N-[2-(3-methoxyphenyl)-ethyl]benzamide (10.1 g, 26 mmol) in CHCl<sub>3</sub> (60 ml) at 0°C under N<sub>2</sub>. The resulting reaction was stirred at 0°C for 10 min, then warmed to rt and stirred at rt for 18 hr. After cooling to 0°C, hexane (500 ml) was added and the resulting suspension was allowed to settle. The solvent was decanted off from the oil that separated and the remaining oil was triturated with EtOH (20 ml) to give a white solid. Et<sub>2</sub>O (200 ml) was added, the resulting suspension was stirred for 1 hr, the solid was collected and dried *in vacuo* to give 7.5 g (72% yield) of white solid.

**Preparation 18****1-(4-Iodophenyl)-6-methoxy-1,2,3,4-tetrahydroisoquinoline**

**[0161]** Sodium borohydride (2 g, 52.9 mmol) was added in small portions to a solution of 1-(4-iodophenyl)-6-methoxy-3,4-dihydroisoquinoline hydrochloride (7.5 g, 18.7 mmol) in MeOH (100 ml) at 0°C. The reaction mixture was stirred at 0°C for 1 hr, then quenched by the dropwise addition of H<sub>2</sub>O (20 ml). The resulting mixture was concentrated *in vacuo* to remove most of the MeOH, then EtOAc was added. The resulting mixture was extracted with 1M NaOH (2 x 25 ml), then saturated NaCl solution (25 ml), dried over MgSO<sub>4</sub>, and evaporated *in vacuo* to give 6.2 g (65% yield) of white solid.

30 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.63 (d, J = 8.52 Hz, 1H), 7.01 (d, J = 8.31 Hz, 2H), 6.66 (d, 1H), 6.61 (d, J = 1.45 Hz, 2H), 5.03 (s, 2H), 4.98 (s, 1H), 3.77 (s, 3H), 3.24 - 3.20 (m, 1H), 3.08 - 2.96 (m, 2H), 2.80 - 2.75 (m, 1H); MS m/e 366 (M<sup>+</sup>+1).

**Preparation 19****Cyclohexanecarboxylic acid [2-(3-methoxyphenyl)ethyl]amide**

**[0162]** A mixture of 3-methoxyphenethylamine (1.000 g, 6.61 mmol), cyclohexanecarboxylic acid (0.847 g, 6.61 mmol), 1-hydroxybenzotriazole hydrate (1.340 g, 9.92 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.330 g, 6.94 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (20 ml) was stirred at 0°C under N<sub>2</sub> for 1 hr, then warmed to rt and stirred at rt for 18 hr. The reaction mixture was washed sequentially with 1M NaOH (10 ml), 1M HCl (10 ml), and H<sub>2</sub>O (10 ml), then dried over MgSO<sub>4</sub> and concentrated *in vacuo* to give 1.301 g of off-white solid.

Purification by flash chromatography, eluting with hexane : EtOAc (6:4) gave 1.107 g (64% yield) of white solid.

45 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.23 - 7.19 (m, 1H), 6.78 - 6.71 (m, 3H), 5.44 (br s, 1H), 3.78 (s, 3H), 3.52 - 3.47 (m, 2H), 2.79 - 2.75 (m, 2H), 1.81 - 1.73 (m, 4H), 1.64 (br s, 1H), 1.41 - 1.33 (m, 2H), 1.26 - 1.18 (m, 4H) MS m/e 262 (M<sup>+</sup>+1).

**Preparation 20****1-Cyclohexyl-6-methoxy-3,4-dihydroisoquinoline**

55 **[0163]** A solution of cyclohexanecarboxylic acid [2-(3-methoxyphenyl)ethyl]amide (1.000 g, 3.8 mmol) in POCl<sub>3</sub> (7.0 ml, 75.1 mmol) was refluxed under N<sub>2</sub> for 24 hr, then the cooled reaction mixture was slowly added to ice-water (30 ml) with stirring. The resulting mixture was extracted with EtOAc (3 x 20 ml), then the remaining aqueous layer was basified to pH 10 with concentrated ammonia (40 ml). This was extracted with CHCl<sub>3</sub> (4 x 20 ml), then the combined

extracts were dried over  $\text{MgSO}_4$  and concentrated *in vacuo* to give 0.270 g (93% yield) of yellow oil.  
 $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.47 (d,  $J$  = 8.72 Hz, 1H), 6.79 (dd,  $J$  = 8.51 Hz,  $J$  = 2.49 Hz, 1H), 6.70 (d,  $J$  = 2.28 Hz, 1H), 3.83 (s, 3H), 3.66 - 3.62 (m, 2H), 2.88 - 2.82 (m, 1H), 2.65 - 2.62 (m, 2H), 1.87 - 1.81 (m, 4H), 1.46 - 1.23 (m, 6H); MS  $m/e$  244 ( $\text{M}^+ + 1$ ).

## Preparation 21

### 1-Cyclohexyl-6-methoxy-1,2,3,4-tetrahydroisoquinoline

[0164] Sodium borohydride (0.512 g, 13.5 mmol) was added in small portions to a solution of 1-cyclohexyl-6-methoxy-3,4-dihydroisoquinoline (0.822 g, 3.38 mmol) in MeOH (40 ml) at  $0^\circ\text{C}$ . The reaction mixture was stirred at  $0^\circ\text{C}$  for 30 min, warmed to rt and stirred at rt for 15 hr, then quenched by the dropwise addition of  $\text{H}_2\text{O}$  (5.0 ml).

The resulting mixture was concentrated *in vacuo* to remove the MeOH and the remaining aqueous solution was extracted with  $\text{CHCl}_3$  (4 x 10 ml). The combined extracts were dried over  $\text{MgSO}_4$  and concentrated *in vacuo* to give 0.775 g (93% yield) of yellow oil.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.05 (d,  $J$  = 8.51 Hz, 1H), 6.73 - 6.70 (dd,  $J$  = 8.72 Hz,  $J$  = 2.91 Hz, 1H), 6.59 (d,  $J$  = 2.70 Hz, 1H), 3.86 (d,  $J$  = 3.94 Hz, 1H), 3.77 (s, 3H), 3.30 - 3.25 (m, 1H), 2.95 - 2.88 (m, 1H), 2.86 - 2.79 (m, 1H), 2.67 - 2.61 (m, 1H), 1.89 - 1.79 (m, 2H), 1.71 - 1.64 (m, 4H), 1.40 - 1.23 (m, 4H), 1.17 - 1.05 (m, 4H); MS  $m/e$  246 ( $\text{M}^+ + 1$ ).

## Preparation 22

### N-[2-(3-Methoxyphenyl)ethyl]benzamide

[0165] Benzoyl chloride (0.922 g, 6.61 mmol) was added to a solution of 3-methoxyphenethylamine (1.000 g, 6.61 mmol) and  $\text{Et}_3\text{N}$  (2.8 ml, 19.8 mmol) in anhydrous THF (30 ml) at rt under  $\text{N}_2$ . The resulting suspension was stirred at rt under  $\text{N}_2$  for 20 hr. The reaction mixture was filtered and the filtrate was concentrated *in vacuo* to give 1.811 g of white solid. Purification by flash chromatography, eluting with hexane : EtOAc (7:3) gave 1.689 g (100% yield) of white solid.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.68 - 7.66 (m, 2H), 7.49 - 7.40 (m, 1H), 7.39 - 7.37 (m, 2H), 7.25 - 7.19 (m, 1H), 6.83 - 6.79 (m, 2H), 6.77 (s, 1H), 6.11 (br s, 1H), 3.78 (s, 3H), 3.74 - 3.69 (m, 2H), 2.92 - 2.89 (m, 2H); MS  $m/e$  256 ( $\text{M}^+ + 1$ ).

## Preparation 23

### 6-Methoxy-1-phenyl-3,4-dihydroisoquinoline

[0166] A solution of N-[2-(3-methoxyphenyl)ethyl]benzamide (1.530 g, 5.99 mmol) in phosphorous oxychloride (10 ml, 107.3 mmol) was refluxed under  $\text{N}_2$  for 24 hours.

The cooled reaction mixture was then slowly added to ice-water (50 ml) with stirring, and basified to pH 10 with concentrated ammonia (60 ml). This was extracted with  $\text{CHCl}_3$  (4 x 25 ml), and the combined organic extracts were dried over  $\text{MgSO}_4$  then concentrated *in vacuo* to give 1.548 g (100% yield) of yellow oil.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.60 - 7.58 (m, 2H), 7.47 - 7.39 (m, 3H), 7.22 (d,  $J$  = 8.51 Hz, 1H), 6.80 (d,  $J$  = 2.70 Hz, 1H), 6.74 (dd,  $J$  = 8.51 Hz,  $J$  = 2.49 Hz, 1H), 3.85 (s, 3H), 3.84 - 3.80 (m, 2H), 2.82 - 2.79 (m, 2H); MS  $m/e$  238 ( $\text{M}^+ + 1$ ).

## Preparation 24

### 6-Methoxy-1-phenyl-1,2,3,4-tetrahydroisoquinoline

[0167] Sodium borohydride (0.945 g, 24.96 mmol) was added in small portions to a solution of 6-methoxy-1-phenyl-3,4-dihydroisoquinoline (1.480 g, 6.24 mmol) in MeOH (50 ml) at  $0^\circ\text{C}$ . The reaction mixture was stirred for 30 min, warmed to rt and stirred at rt for 19 hr, then quenched by the dropwise addition of  $\text{H}_2\text{O}$  (10 ml). Excess MeOH was removed by evaporation and the resulting aqueous solution was extracted with  $\text{CHCl}_3$  (4 x 10 ml). The combined organic extracts were dried over  $\text{MgSO}_4$  and concentrated *in vacuo* to give 1.308 g (88% yield) of white solid.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.33 - 7.26 (m, 5H), 6.66 - 6.59 (m, 3H), 5.05 (s, 1H), 3.76 (s, 3H), 3.26 - 3.20 (m, 1H), 3.08 - 2.99 (m, 2H), 2.84 - 2.77 (m, 1H); MS  $m/e$  240 ( $\text{M}^+ + 1$ ).

**Preparation 25****Thiophene-2-carboxylic acid [2-(3-methoxyphenyl)ethyl]amide**

**[0168]** A mixture of 3-methoxyphenethylamine (1.000 g, 6.61 mmol), 2-thiophenecarboxylic acid (0.847 g, 6.61 mmol), 1-hydroxybenzotriazole hydrate (1.340 g, 9.92 mmol), and 1-(3-diethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.330 g, 6.94 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (20 ml) was stirred at 0°C under  $\text{N}_2$  for 1 hr, then warmed to rt and stirred at rt for 18 hr. The reaction mixture was washed with 1M NaOH (10 ml) followed by 1M HCl (10 ml), dried over  $\text{MgSO}_4$ , and concentrated *in vacuo* to give 1.859 g of yellow oil. Purification by flash chromatography, eluting with hexane : EtOAc (7:3) gave 1.637 g (95% yield) of yellow oil.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.43 (dd,  $J$  = 4.98 Hz,  $J$  = 1.04 Hz, 1H), 7.38 (dd,  $J$  = 3.74 Hz,  $J$  = 1.25 Hz, 1H), 7.25 - 7.21 (m, 1H), 7.03 (dd,  $J$  = 4.98 Hz,  $J$  = 3.74 Hz, 1H), 6.82 - 6.76 (m, 3H), 5.97 (br s, 1H), 3.78 (s, 3H), 3.70 - 3.65 (m, 2H), 2.90 - 2.87 (m, 2H); MS  $m/e$  262 ( $\text{M}^+ + 1$ ).

**Preparation 26****6-Methoxy-1-thiophen-2-yl-3,4-dihydroisoquinoline**

**[0169]** A solution of thiophene-2-carboxylic acid [2-(3-methoxyphenyl)ethyl]amide (1.480 g, 5.66 mmol) in phosphorous oxychloride (10 ml, 107.3 mmol) was refluxed under  $\text{N}_2$  for 24 hr. The cooled reaction mixture was then slowly added to ice-water (50 ml) with stirring and basified to pH 10 with concentrated ammonia (60 ml). This was extracted with  $\text{CHCl}_3$  (4 x 25 ml) and the combined extracts were dried over  $\text{MgSO}_4$  then concentrated *in vacuo* to give 1.517 g (96% yield) of yellow oil.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.66 (d,  $J$  = 8.51 Hz, 1H), 7.48 - 7.45 (m, 2H), 7.13 - 7.11 (m, 1H), 6.88 - 6.81 (m, 2H), 3.87 (s, 3H), 3.80 - 3.76 (m, 2H), 2.79 - 2.75 (m, 2H); MS  $m/e$  244 ( $\text{M}^+ + 1$ ).

**Preparation 27****6-Methoxy-1-thiophen-2-yl-1,2,3,4-tetrahydroisoquinoline**

**[0170]** Sodium borohydride (0.916 g, 24.2 mmol) was added in small portions to a solution of 6-methoxy-1-thiophen-2-yl-3,4-dihydroisoquinoline (1.470 g, 6.04 mmol) in MeOH (50 ml) at 0°C. The reaction mixture was stirred for 30 min, warmed to rt and stirred at rt for 21 hr, then quenched by the dropwise addition of  $\text{H}_2\text{O}$  (10 ml). Excess MeOH was removed by evaporation and the resulting aqueous solution was extracted with  $\text{CHCl}_3$  (4 x 10 ml). The combined extracts were dried over  $\text{MgSO}_4$  and concentrated *in vacuo* to give 1.316 g (89% yield) of yellow solid.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.23 (d,  $J$  = 2.91 Hz, 1H), 6.94 - 6.88 (m, 3H), 6.68 - 6.65 (m, 2H), 5.39 (s, 1H), 3.77 (s, 3H), 3.28 - 3.22 (m, 1H), 3.11 - 3.05 (m, 1H), 2.97 - 2.90 (m, 1H), 2.87 - 2.82 (m, 1H); MS  $m/e$  246 ( $\text{M}^+ + 1$ ).

**Preparation 28****4-Bromo-N-[2-(3-methoxyphenyl)ethyl]benzamide**

**[0171]** To a stirred solution of 3-methoxyphenethylamine (25.0 g, 165 mmol) and  $\text{Et}_3\text{N}$  (30.0 ml, 214.50 mmol) in  $\text{CH}_2\text{Cl}_2$  (500 ml) at 0°C was added 4-bromobenzoyl chloride (25 g, 181 mmol) in portions, and the mixture stirred for 30 min. 2N HCl was added and stirred for 5 min, and the aqueous layer was separated. The organic phase was washed sequentially with 1N HCl (1 x 200 ml),  $\text{H}_2\text{O}$  (2 x 100 ml), and saturated  $\text{NaHCO}_3$  solution (100 ml), dried over  $\text{MgSO}_4$ , and concentrated *in vacuo* to give 28.42 g (52% yield) of off-white solid.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.53 (m, 4H), 7.23 (m, 1H), 6.82-6.73 (m, 3H), 3.77 (s, 3H), 3.75-3.69 (m, 2H), 2.89 (t, 2H); MS  $m/e$  334, 336 bromine isotope pattern ( $\text{M}^+ + 1$ ).

**Preparation 29****1-(4-Bromophenyl)-6-methoxy-3,4-dihydroisoquinoline hydrochloride**

**[0172]** To a solution of 4-bromo-N-[2-(3-methoxyphenyl)ethyl]benzamide (5.00 g, 15.0 mmol) in  $\text{CHCl}_3$  (30 ml) was added phosphorous pentachloride (5.30 g, 25.0 mmol) and the mixture stirred for 18 hr at rt under  $\text{N}_2$ . Hexane was added to the mixture and the liquid was then decanted off. EtOH (15 ml) was added to the residue and the mixture was triturated with diethyl ether to give 3.65 g (69% yield) of off-white solid.

<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 7.86 (d, J = 8.51 Hz, 2H), 7.61 (d, J = 8.30 Hz, 2H), 7.47 (d, J = 8.92 Hz, 1H), 7.04 (d, J = 2.49 Hz, 1H), 7.04 (dd, J = 8.92 Hz, J = 2.70 Hz, 1H), 3.98-3.94 (m, 5H), 3.29-3.23 (m, 2H).

### Preparation 30

#### 1-(4-Bromophenyl)-6-methoxy-1,2,3,4-tetrahydroisoquinoline

[0173] Sodium borohydride (0.720g, 19.0 mmol) was added in small portions to a solution of 1-(4-bromophenyl)-6-methoxy-3,4-dihydroisoquinoline, hydrochloride (3.62 g, 10.2 mmol) in MeOH (30 ml) at 0°C. The reaction mixture was stirred for 30 min, warmed to rt, and stirred at rt for 1 hr, then quenched by the dropwise addition of H<sub>2</sub>O (10 ml). Acetic acid (10 drops) was added with stirring, and the resulting mixture was basified to pH 10 with 1N NaOH and extracted with EtOAc (3 x 25 ml). The combined extracts were washed with 1N NaOH followed by saturated NaCl solution, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated *in vacuo* to give 3.24 g (100% yield) of white solid.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.43 (d, J = 8.51 Hz, 1H), 7.15 (d, J = 8.51 Hz, 1H), 6.66 - 6.60 (m, 3H), 5.02 (s, 1H), 3.76 (s, 3H), 3.23 - 3.17 (m, 1H), 3.07 - 3.01 (m, 2H), 2.84 - 2.78 (m, 1H).

### Preparation 31

#### 2-Benzenesulfonyl-6-benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinoline

[0174] Benzenesulfonyl chloride (0.016 g, 0.09 mmol) was added to a solution of 6-benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinoline (0.040 g, 0.09 mmol) and Et<sub>3</sub>N (0.026 ml, 0.19 mmol) in anhydrous THF (10 ml) at rt under N<sub>2</sub>. The resulting suspension was stirred at rt for 18 hr, then evaporated *in vacuo* to give 0.086 g of yellow residue. Purification by flash chromatography, eluting with EtOAc: MeOH (6:4) gave 0.045 g (85% yield) of white solid.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.65 (d, J = 7.26 Hz, 3H), 7.38 (s, 5H), 7.34 - 7.32 (m, 1H), 7.28 (d, J = 7.06 Hz, 3H), 7.07 (d, J = 7.47 Hz, 3H), 6.86 (d, J = 7.47 Hz, 1H), 6.76 (d, J = 6.64 Hz, 4H), 6.54 (s, 1H), 6.14 (s, 1H), 4.98 (s, 2H), 4.28- 4.24 (m, 1H), 3.78 - 3.74 (m, 1H), 3.33 - 3.24 (m, 4H), 2.55 - 2.45 (m, 2H), 1.96 (br s, 4H); MS m/e 569 (M<sup>+</sup>+1).

### Preparation 32

#### 6-Benzyloxy-2-(naphthalene-1-sulfonyl)-1-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-1,2,3,4-tetrahydroisoquinoline

[0175] A solution of 6-benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinoline (0.015 g, 0.04 mmol) and Et<sub>3</sub>N (0.01 ml, 0.07 mmol) in anhydrous THF (0.35 ml) was added to a solution of 1-naphthalenesulfonyl chloride (0.04 mmol) in anhydrous THF (0.4 ml) at rt in a sealed reaction vessel. The resulting suspension was stirred at rt for 20 hr, then evaporated to dryness. The product was suspended in a mixture of H<sub>2</sub>O (0.4 ml) and saturated NaHCO<sub>3</sub> solution (0.4 ml), then extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 0.75 ml). The combined extracts were evaporated to dryness to give the crude product. Purification by reverse-phase HPLC on a Primesphere C-18HC (50.0 mm x 10.0 mm column with 5 μm particle size) column, eluting with a linear gradient starting at time 0 min. of H<sub>2</sub>O : CH<sub>3</sub>CN : 1%TFA/H<sub>2</sub>O (85 : 10 : 5), increasing to H<sub>2</sub>O : CH<sub>3</sub>CN : 1%TFA/H<sub>2</sub>O (5 : 90 : 5) at 8 min., detected on a Micromass Platform 2 mass spectrometer (DAD 190 - 600 nm) gave material which, after evaporation to dryness, was purified by reverse-phase HPLC on a Primesphere C-18HC (3.0 mm x 2.0 mm column with 5 μm particle size) column, eluting with a linear gradient starting at time 0 min. of H<sub>2</sub>O : CH<sub>3</sub>CN : TFA (99.9 : 0 : 0.1), increasing to H<sub>2</sub>O : CH<sub>3</sub>CN : TFA (0 : 99.9 : 0.1) at 4 min., detected with a UV detector (300 nm +/- 90 nm and 254 nm +/- 25 nm), gave an eluent that was evaporated to dryness to give the named product. MS m/e 583 (M<sup>+</sup>+1).

### Preparation 33

#### 6-Benzyloxy-2-(3,5-dimethylisoxazole-4-sulfonyl)-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinoline

[0176] 3,5-Dimethylisoxazole-4-sulfonyl chloride (0.034 g, 0.18 mmol) was added to a solution of 6-benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinoline (0.075 g, 0.18 mmol) and Et<sub>3</sub>N (0.037 ml, 0.26 mmol) in anhydrous THF (10 ml) at rt under N<sub>2</sub>. The resulting suspension was stirred at rt for 21 hr, then evaporated *in vacuo* to a residue that was suspended in a mixture of saturated NaHCO<sub>3</sub> solution (4 ml) and H<sub>2</sub>O (4 ml). This was extracted with CHCl<sub>3</sub> (4 x 5 ml) and the combined extracts were dried over MgSO<sub>4</sub>, then concentrated *in vacuo* to give 0.110 g of yellow oil. Purification by flash chromatography, eluting with EtOAc : MeOH (1:1) gave 0.078 g (76% yield) of white

solid.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.42 - 7.37 (m, 2H), 7.33 (d, J = 6.65 Hz, 1H), 7.19 - 7.10 (m, 1H), 7.02 - 6.98 (m, 2H), 6.90 - 6.87 (m, 2H), 6.81 - 6.79 (m, 3H), 6.69 (s, 1H), 6.05 (s, 1H), 5.01 (s, 2H), 4.08 - 4.03 (m, 2H), 3.72 - 3.67 (m, 1H), 3.35 - 3.30 (m, 1H), 2.87 - 2.84 (m, 2H), 2.74 - 2.71 (m, 1H), 2.67 - 2.66 (m, 1H), 2.59 (s, 4H), 2.55 (s, 1H), 2.52 (s, 2H), 2.19 (s, 2H), 2.17 (s, 1H), 1.78 (br s, 4H); MS m/e 588 (M<sup>+</sup>+1).

#### Preparation 34

##### 1-[4-(2-Pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol

**[0177]** A mixture of 6-benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinoline (0.300 g, 0.70 mmol), ammonium formate (0.883 g, 14.0 mmol), and 20% Pd(OH)<sub>2</sub>/C (0.050 g) in MeOH (50 ml) was refluxed under N<sub>2</sub> for 1 hr, then filtered through Celite®. The filtrate was evaporated *in vacuo* to a yellow residue that was suspended in a mixture of H<sub>2</sub>O (5 ml) and saturated NaHCO<sub>3</sub> solution (5 ml), then extracted with CH<sub>2</sub>Cl<sub>2</sub> (4 x 10 ml). The combined organic extracts were dried over MgSO<sub>4</sub> and concentrated *in vacuo* to give 0.133 g of yellow oil. The aqueous layer was extracted again with CH<sub>2</sub>Cl<sub>2</sub> (4 x 50 ml) to give an additional 0.083 g of yellow oil, for a total of 0.216 g (91% yield) of product.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.54 (d, J = 8.51 Hz, 1H), 7.09 (d, J = 8.51 Hz, 1H), 6.83 (d, J = 8.72 Hz, 1H), 6.72 (d, J = 8.51 Hz, 1H), 6.57 - 6.47 (m, 3H), 4.97 (s, 1H), 4.09-4.05 (m, 2H), 3.18 - 3.15 (m, 1H), 3.01 - 2.88 (m, 4H), 2.80 - 2.79 (m, 1H), 2.69 (br s, 4H), 2.16 - 2.15 (m, 1H), 1.83 (br s, 4H); MS m/e 339 (M<sup>+</sup>+1).

#### Preparation 35

##### 6-Benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinoline-2-carboxylic acid tert-butyl ester

**[0178]** A solution of di-tert-butyl dicarbonate (0.035 g, 0.16 mmol), 6-benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinoline (0.069 g, 0.16 mmol), and Et<sub>3</sub>N (0.028 ml, 0.20 mmol) in anhydrous THF (10 ml) was stirred at rt under N<sub>2</sub> for 20 hr. The reaction mixture was concentrated *in vacuo* to give 0.081 g of yellow oil. Purification by flash chromatography, eluting with EtOAc : MeOH (6:4), gave 0.041 g (48% yield) of white solid.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.43 - 7.27 (m, 5H), 7.20 - 7.14 (m, 1H), 7.10 (d, J = 8.72 Hz, 2H), 6.90 (br s, 1H), 6.80 - 6.77 (m, 4H), 5.03 (s, 2H), 4.18 (br s, 2H), 3.13 - 2.65 (m, 10H), 1.89 (br s, 4H), 1.47 (s, 9H); MS, m/e 529 (M<sup>+</sup>+1).

#### Preparation 36

##### 6-Benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinoline-2-carboxylic acid ethyl ester

**[0179]** A solution of ethyl chloroformate (0.017 g, 0.16 mmol), 6-benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinoline (0.069 g, 0.16 mmol), and Et<sub>3</sub>N (0.028 ml, 0.20 mmol) in anhydrous THF (10 ml) was stirred at rt under N<sub>2</sub> for 20 hr. The reaction mixture was concentrated *in vacuo* to give 0.115 g of tan solid.

Purification by flash chromatography, eluting with EtOAc : MeOH (8:2), gave 0.023 g (28% yield) of white solid.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.43 - 7.29 (m, 6H), 7.12 - 7.10 (m, 2H), 6.91 (d, J = 7.68 Hz, 1H), 6.80 - 6.77 (m, 4H), 5.04 (s, 2H), 4.29 - 3.98 (m, 7H), 3.25 - 2.67 (m, 7H), 1.97 (br s, 4H), 1.30 - 1.23 (s, 3H); MS, m/e 501 (M<sup>+</sup>+1).

#### Preparation 37

##### 2-(3-Methoxy-phenyl)-N-phenyl-acetamide

**[0180]** The title compound was prepared according to the procedure of Nagarajan et al., *Ind. J. Chem.*, 24B:83-97 (1985).

#### Preparation 38

##### [2-(3-Methoxy-phenyl)-ethyl]-phenyl-amine

**[0181]** The title compound was prepared according to the procedure of Nagarajan et al., *Ind. J. Chem.*, 24B:83-97 (1985).

**Preparation 39****4-Methoxy-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide**

**[0182]** To a stirred solution of 2-(3-methoxy-phenyl)-ethyl]-phenyl-amine (0.800g, 3.52 mmol, 1.0 eq.) in CH<sub>2</sub>Cl<sub>2</sub> (4ml) was added Et<sub>3</sub>N (2.97 ml, 21.1 mmol, 6.0 eq.), PPAA (50% solution in EtOAc, 7.05 mmol, 2.0 eq.), para-methoxy benzoic acid (0.803g, 5.29 mmol, 1.5 eq.), and a catalytic amount of DMAP. The reaction was stirred at room temperature for 36 hours. The reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> (30 ml) and washed successively with 1N HCl (1x20 ml) and saturated NaHCO<sub>3</sub> (1x20 ml). The organic layer was dried over MgSO<sub>4</sub>, filtered, and concentrated in *vacuo*. Flash chromatography (SiO<sub>2</sub>, hexanes : EtOAc 7:1 to 3:1) of the residue afforded the desired compound as a colorless oil (1.13g, 3.13 mmol, 89% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 2.95 (2H, t, J 8.0 Hz) 3.68 (3H, s), 3.73 (3H,s), 4.08 (2H, t, J 8.0 Hz), 6.60-7.24 (13H, overlapping m).

**Preparation 40****Cyclohexanecarboxylic acid [2-(3-methoxy-phenyl)-ethyl]-phenyl-amide**

**[0183]** The title compound was prepared by analogy to Preparation 39 except that cyclohexane carboxylic acid was used instead of *para*-methoxy benzoic acid, and hexanes 7:1 EtOAc was used as the eluent for flash chromatography purification. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 0.84-0.94 (2H, overlapping m), 1.09-1.23 (2H, overlapping m), 1.44-1.77 (6H, overlapping m), 2.05 (1H, m), 2.81 (2H, m), 3.73 (3H, s), 3.85 (2H, m), 6.68-6.73 (3H, overlapping m), 7.00-7.02 (2H, overlapping m) 7.13 (1H, m) and 7.31-7.38 (3H overlapping m). MS 338 (M+1).

**Preparation 41****N-[2-(3-Methoxy-phenyl)-ethyl]-N-phenyl-isobutyramide**

**[0184]** The title compound was prepared by analogy to Preparation 39 except that 2-methyl propanoic acid was used instead of *para*-methoxy benzoic acid, and hexanes 7:1 EtOAc was used as eluent for flash chromatography purification. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 1.00 (6H, d, J 6.5 Hz), 2.40 (1H, septet, J 6.5 Hz), 2.85 (2H, t, 7.5 Hz), 3.75 (3H, s), 3.88 (2H, t, 7.5 Hz) 6.72-6.73 (3H, overlapping m) and 7.31-7.41 (9H overlapping m). MS 298 (M+1).

**Preparation 42****N-[2-(3-Methoxy-phenyl)-ethyl]-N-phenyl-benzamide**

**[0185]** To a stirred solution of 2-(3-methoxy-phenyl)-ethyl]-phenyl-amine (1.38g, 6.08 mmol, 1.0 eq.) in CH<sub>2</sub>Cl<sub>2</sub> (30 ml) under an atmosphere of N<sub>2</sub> was added EDC (1.05g, 6.69 mmol, 1.1 eq.) and HOBt (1.23g, 91.2 mmol, 1.5 eq.). Benzoic acid (1.11g, 9.12 mmol, 1.5 eq.) was added to this mixture, and stirring was continued at room temperature for 24 hours. Additional EDC (1.1g, 7.01 mmol, 1.15 eq.) and a catalytic amount of DMAP were added, and stirring was continued at room temperature for a further 24 hours. The reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> (30 ml) and washed sequentially with sat. NaHCO<sub>3</sub> (1x30 ml) and 1N HCl (1x30 ml). The basic aqueous layer was back extracted with CH<sub>2</sub>Cl<sub>2</sub> (1x20 ml). The combined organics were dried over MgSO<sub>4</sub>, filtered, and concentrated in *vacuo*. The residue was subjected to flash chromatography (SiO<sub>2</sub>, hexanes:EtOAc 8:1) to give the desired product (1.41g, 4.24 mmol, 70% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 2.96 (2H, t, J 8.0 Hz), 3.74 (3H, s), 4.08 (2H, t, J 8.0 Hz) and 6.73-7.26 (14H, overlapping m). MS 332 (M+1).

**Preparation 43****4-(2E-Ethoxycarbonyl-vinyl)-benzoic acid**

**[0186]** To a stirred solution of 4-formyl-benzoic acid (10.0g, 66.7 mmol) in THF (140 ml) was added carboethoxyl-riphenylphosphorane (25.0g, 71.8 mmol). To this solution was added NaOH (2.50g, 66.7mmol) as a solution in water (20 ml). The reaction was stirred at rt overnight. The reaction mixture was diluted with water (50ml) and extracted with

EtOAc (3x50 ml). The combined extracts were dried over  $\text{MgSO}_4$ , filtered, and concentrated in *vacuo*. The residue was purified by flash chromatography ( $\text{SiO}_2$ , 5%  $\text{MeOH}/\text{CH}_2\text{Cl}_2$ ) to give the desired compound as a white solid (11.68g, 53.1 mmol, 80% yield).

$^1\text{H}$  NMR (400 MHz,  $\text{d}_6$ -acetone)  $\delta_{\text{H}}$  1.28 (3H, t, J 7.0 Hz), 4.20 (2H, q, J 7.0 Hz), 6.65 (1H, d, J 16.0 Hz) and 7.67-8.13 (5H, overlapping m).

#### Preparation 44

##### 3-(4-[[2E-(3-Methoxy-phenyl)-ethyl]-phenyl-carbamoyl]-phenyl)-acrylic acid ethyl ester

**[0187]** The title compound was prepared by analogy to Preparation 39 except that 4-(2E-ethoxycarbonyl-vinyl)-benzoic acid was used instead of *para*-methoxy benzoic acid and the crude material was taken to the next step without any flash chromatography.

MS 430 (M+1).

#### Preparation 45

##### N-[2-(3-Methoxy-phenyl)-ethyl]-2,N-diphenyl-acetamide

**[0188]** The title compound was prepared by analogy to Preparation 39 except that phenyl acetic acid was used instead of *para*-methoxy benzoic acid, and the crude material was taken to the next step without any flash chromatography.

MS 346 (M+1).

#### Preparation 46

##### Thiophene-2-carboxylic acid [2-(3-methoxy-phenyl)-ethyl]-phenyl-amide

**[0189]** The title compound was prepared by analogy to Preparation 39 except that thiophene-2-carboxylic acid was used instead of *para*-methoxy benzoic acid, and hexanes 10:1 EtOAc was used as eluent for flash chromatography purification.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  2.98 (2H, m), 3.75 (3H, s), 4.05 (2H, m), 6.65-6.80 (5H, overlapping m) and 7.11-7.39 (7H, overlapping m). MS 337 (M).

#### Preparation 47

##### Naphthalene-2-carboxylic acid [2-(3-methoxy-phenyl)-ethyl]-phenyl-amide

**[0190]** The title compound was prepared by analogy to Preparation 39 except that naphthalene-2-carboxylic acid was used instead of *para*-methoxy benzoic acid, and the crude material was taken to the next step without any flash chromatography.

MS 382 (M+1).

#### Preparation 48

##### 3,4,5-Trifluoro-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide

**[0191]** The title compound was prepared by analogy to Preparation 39 except that 3,4,5-trifluorobenzoic acid was used instead of *para*-methoxy benzoic acid, and the crude material was taken to the next step without any flash chromatography.

MS 386 (M+1).

#### Preparation 49

##### 4-Chloro-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide

**[0192]** The title compound was prepared by analogy to Preparation 39 except that 4-chloro-benzoic acid was used instead of *para*-methoxy benzoic acid, and the crude material was taken to the next step without any flash chromatography.



raphy.  
MS 365 (M+1).

#### Preparation 50

##### Thiazole-2-carboxylic acid [2-(3-methoxy-phenyl)-ethyl]-phenyl-amide

[0193] The title compound was prepared by analogy to Preparation 39 except that thiazole-2-carboxylic acid (see Metzger, *Bull. Soc. Chim. Fr.*, p. 708 (1953)) was used instead of *para*-methoxy benzoic acid, and the crude material was taken to the next step without any flash chromatography.  
MS 338 (M+1).

#### Preparation 51

##### Adamantane-1-carboxylic acid [2-(3-methoxy-phenyl)-ethyl]-phenyl-amide

[0194] The title compound was prepared by analogy to Preparation 39 except that adamantane-1-carboxylic acid was used instead of *para*-methoxy benzoic acid, and the crude material was taken to the next step without any flash chromatography.  
MS 389 (M).

#### Preparation 52

##### N-[2-(3-Methoxy-phenyl)-ethyl]-N-phenyl-isonicotinamide

[0195] The title compound was prepared by analogy to Preparation 39 except that isonicotinic acid was used instead of *para*-methoxy benzoic acid, and hexanes:EtOAc 4:1 was used as eluent for flash chromatography purification.  
<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 2.95 (2H, m), 3.75 (3H, s), 4.10 (2H, m), 6.74-6.85 (3H, overlapping m), 7.08-7.23 (8H, overlapping m) and 8.41 (2H, m). MS 333 (M+1).

#### Preparation 53

##### 4-Iodo-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide

[0196] To a stirred suspension of 4-iodobenzoic acid (8.5g, 34.4 mmol, 1.3 eq.) in toluene (21ml) under an atmosphere of N<sub>2</sub> was added thionyl chloride (17ml). The suspension was heated at reflux for 2 hours forming a turbid solution. The reaction was allowed to cool to RT and the volatiles were removed in *vacuo*. The residue was taken up in THF (100 ml) under an atmosphere of N<sub>2</sub> and cooled to 0°C. To the ice cold solution was added successively Et<sub>3</sub>N (14.7 ml, 106.0 mmol, 4.0 eq.), a catalytic amount of DMAP, and 2-(3-methoxy-phenyl)-ethyl-phenyl-amine (6.0 g, 26.4 mmol, 1.0 eq.). After 1 hour at 0°C the ice bath was removed and stirring was continued at RT overnight. The reaction mixture was diluted with EtOAc (100ml) and washed successively with 1N HCl (2x100ml), water (2x100ml), and 10% K<sub>2</sub>CO<sub>3</sub> (2x100ml). The organic layer was dried over MgSO<sub>4</sub>, filtered, and concentrated in *vacuo*. The residue was subjected to flash chromatography (SiO<sub>2</sub>, hexanes:EtOAc 8:1 to 4:1 gradient) to give the desired product (11.0 g, 24.1 mmol, 91% yield).  
<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 2.96 (2H, t, J 8.0 Hz), 3.77 (3H, s), 4.10 (2H, t, J 8.0 Hz), 6.75-6.87 (5H, overlapping m), 6.99 (2H, d, J 8.5 Hz), 7.15-7.24 (4H, overlapping m) and 7.49 (2H, d, J 8.5 Hz). MS 458 (M+1).

#### Preparation 54

##### 1-Trifluoroacetyl-piperidine-4-carbonyl chloride

[0197] The title compound was prepared according to the procedure of Hibert et al., *J. Med. Chem.*, 33:1594 (1990).  
**[Was this done exactly as described, or analogously? Done exactly]**

**Preparation 55****1-Trifluoroacetyl-piperidine-4-carboxylic acid [2-(3-methoxy-phenyl)-ethyl]-amide**

[0198] To a stirred solution of 3-methoxy phenylethylamine (9.3g, 61.6 mmol) and triethylamine (8.1 g, 11.2 ml, 80.1 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (50 ml) at 0°C under an atmosphere of N<sub>2</sub> was added 1-trifluoroacetyl-piperidine-4-carboxyl chloride (15.0 g, 61.6 mmol) as a solution in CH<sub>2</sub>Cl<sub>2</sub> (50 ml) in a dropwise manner. The reaction was allowed to warm to RT overnight. The reaction was quenched with water (50 ml), the layers were separated, and the organic layer was washed with water (1x50 ml). The combined aqueous layers were back extracted with CH<sub>2</sub>Cl<sub>2</sub> (1x75 ml). The combined organic layers were washed with brine (1x75 ml). The organic layer was dried over MgSO<sub>4</sub>, filtered, and concentrated in *vacuo*. The resulting solid was triturated with Et<sub>2</sub>O and filtered to give the desired product (18.8 g, 52.3 mmol, 85% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 1.58-1.97 (5H, overlapping m), 2.30 (1H, m), 2.79 (2H, t, J 6.5 Hz), 2.91 (1H, m), 3.17 (1H, m), 3.52 (2H, m), 3.79 (3H, s), 3.99 (1H, m), 4.41 (1H, m), 5.45 (1H, broad s), 6.71-6.79 (3H, overlapping m) and 7.23 (1H, t, J 8.0 Hz). MS 359 (M+1).

**Preparation 56****2,2,2-Trifluoro-1-[4-(6-methoxy-3,4-dihydro-isoquinolin-1-yl)-piperidin-1-yl]-ethanone**

[0199] A stirred solution of 1-trifluoroacetyl-piperidine-4-carboxylic acid [2-(3-methoxy-phenyl)-ethyl]-amide (1.04g, 2.9 mmol) in POCl<sub>3</sub> under an atmosphere of N<sub>2</sub> was heated at reflux for 2.5 hours. The reaction was allowed to cool to RT and the POCl<sub>3</sub> was removed in *vacuo*. The residual oil was suspended in toluene (20 ml) and concentrated in *vacuo* (this process was repeated twice more), leaving the desired product as a tan colored solid (0.872g, 2.56 mmol, 88% yield). MS 341 (M+1)

**Preparation 57****6-Methoxy-1-piperidin-4-yl-3,4-dihydro-isoquinoline**

[0200] To a stirred solution of 2,2,2-trifluoro-1-[4-(6-methoxy-3,4-dihydro-isoquinolin-1-yl)-piperidin-1-yl]-ethanone (0.160g, 0.470 mmol) was added a solution of 10% K<sub>2</sub>CO<sub>3</sub> (2 ml). The cloudy mixture was stirred at RT overnight. The reaction mixture was extracted with EtOAc (3x5 ml), the combined organic extracts were washed with brine (1x5 ml), dried over MgSO<sub>4</sub>, filtered, and concentrated in *vacuo* to give the desired compound as a yellow oil (0.101 g, 0.413 mmol, 88% yield). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 1.67-1.81 (2H, overlapping m), 1.83-2.02 (2H overlapping m), 2.60 (2H, t, J 7.5 Hz), 2.86 (2H, m), 3.08 (1H, m), 3.28 (2H, m), 3.61 (2H, t, J 7.5 Hz), 3.81 (3H, s), 5.24 (1H, broad s), 6.70 (1H, d, J 3.0 Hz), 6.78 (1H, dd, J 8.0 and 3.0 Hz) and 7.38 (1H, d, J 8.0 Hz). MS 245 (M+1).

**Preparation 58****6-Methoxy-1-[1-(1-methyl-1H-imidazole-4-sulfonyl)-piperidin-4-yl]-3,4-dihydro-isoquinoline**

[0201] To a stirred solution of 6-methoxy-1-piperidin-4-yl-3,4-dihydro-isoquinoline (0.094 g, 0.385 mmol) and triethylamine (0.078 g, 0.77 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 ml) was added 1-methyl-1H-imidazole-4-sulfonyl chloride (0.070 g, 0.385 mmol). The reaction mixture was stirred at RT under an atmosphere of N<sub>2</sub> overnight. The reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> (10 ml) and washed with water (2x5 ml). The organic layer was dried over MgSO<sub>4</sub>, filtered, and concentrated in *vacuo*. The residue was purified via radial chromatography (SiO<sub>2</sub>, 1 mm, CH<sub>2</sub>Cl<sub>2</sub> to 5% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to give the product as an oil (0.084 g, 0.216 mmol, 56% yield). MS 389 (M+1).

**Preparation 59****6-Methoxy-1-[1-(1-methyl-1H-imidazole-4-sulfonyl)-piperidin-4-yl]-1,2,3,4-tetrahydroisoquinoline**

[0202] To a stirred solution of 6-methoxy-1-[1-(1-methyl-1H-imidazole-4-sulfonyl)-piperidin-4-yl]-3,4-dihydro-isoquinoline (0.077 g, 0.197 mmol) in MeOH (3 ml) cooled to 0°C was added NaBH<sub>4</sub>. The reaction mixture was then stirred at RT for 2 hours and diluted with water (5 ml) and sat. NaHCO<sub>3</sub> (5 ml). The mixture was extracted with EtOAc (3x10

ml). The combined organics were washed with brine (1x10 ml), dried over  $\text{MgSO}_4$ , filtered, and concentrated *in vacuo*. The residue was purified by radial chromatography ( $\text{SiO}_2$ , 1 mm,  $\text{CH}_2\text{Cl}_2$  to 5%  $\text{MeOH}/\text{CH}_2\text{Cl}_2$ ) to give the product as an oil (0.055 g, 0.141 mmol, 71% yield).  
MS 391 (M+1).

## Preparation 60

### 2-Benzylamino-1-(4-methoxy-phenyl)-ethanone

**[0203]** To a solution of  $\text{Et}_3\text{N}$  (13.55 g, 133.9 mmol) and benzylamine (11.96 g, 111.6 mmol) in THF (25 ml) was added 2-bromo-1-(4-methoxy-phenyl)-ethanone (25.56 g, 111.6 mmol). The reaction was stirred at RT for 60 minutes, then filtered and the filtrate concentrated *in vacuo*. The residue was purified via flash chromatography ( $\text{SiO}_2$ , gradient column 35% to 80%  $\text{EtOAc}/\text{hexanes}$ ) to give the title product (17.12 g, 67.0 mmol, 60% yield).

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  3.85 (3H, s), 3.94 (s, 2H), 4.11 (s, 2H), 6.91 (2H, d,  $J = 9.0$  Hz), 7.20-7.41 (m, 5H) and 7.87 (2H, d,  $J$  9.0 Hz). MS 256 (M+1).

## Preparation 61

### 2-Benzylamino-1-(4-methoxy-phenyl)-ethanol

**[0204]** To a solution of 2-benzylamino-1-(4-methoxy-phenyl)-ethanone (1.61 g, 6.30 mmol, see Preparation 60) in MeOH (60 ml) was added  $\text{NaBH}_4$  (0.48g, 12.6 mmol) in three equal portions. The reaction was stirred at RT overnight then it was quenched with a 1:1 mixture of water and sat.  $\text{NaHCO}_3$  (30 ml). The mixture was extracted with  $\text{CH}_2\text{Cl}_2$  (3x40 ml), the organics were combined, dried over  $\text{MgSO}_4$ , filtered, and concentrated *in vacuo*. The residue was purified via Biotage flash chromatography ( $\text{SiO}_2$ , neat  $\text{EtOAc}$ ) to give the title product (1.216 g, 4.72 mmol, 75% yield).

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  2.68 (2H, broad s), 2.71 (1H, dd,  $J$  12.5 and 9.0 Hz), 2.87 (1H, dd,  $J$  12.5 and 3.5 Hz), 3.77 (3H, s), 3.81 (2H, m), 4.67 (1H, dd  $J$  9.0 Hz and 3.5 Hz), 6.85 (2H, d  $J$  8.5 Hz) and 7.19-7.38 (7H, m). MS 258 (M+1).

## Preparation 62

### 2-[Benzyl-(3-methoxy-benzyl)-amino]-1-(4-methoxy-phenyl)-ethanone

**[0205]** To a solution of 2-benzylamino-1-(4-methoxy-phenyl)-ethanone (12.0 g, 47.0 mmol, see Preparation 60) and 3-methoxy benzaldehyde (6.09 g, 44.8 mmol) in 1,2 DCE (250 ml) was added  $\text{NaB}(\text{OAc})_3\text{H}$ . The reaction mixture was stirred at RT overnight. The reaction was poured into sat.  $\text{NaHCO}_3$  (150 ml) and extracted with  $\text{CH}_2\text{Cl}_2$  (3x 100 ml). The combined organics were dried over  $\text{MgSO}_4$ , filtered, and concentrated *in vacuo*. Flash chromatography ( $\text{SiO}_2$ , gradient elution 20% to 50%  $\text{EtOAc}/\text{hexanes}$ ) gave the title product (13.85 g, 36.9 mmol, 83% yield).

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  3.76 (2H, s), 3.77 (3H, s), 3.78 (2H, s), 3.83 (2H, s), 3.84 (3H, s), 6.83-6.93 (5H, m), 7.19-7.36 (6H, m) and 7.83 (2H, d,  $J$  7.5 Hz). MS 376 (M+1).

## Preparation 63

### 2-[Benzyl-(3-methoxy-benzyl)-amino]-1-(4-methoxy-phenyl)-ethanol

**[0206]** To a solution of 2-[benzyl-(3-methoxy-benzyl)-amino]-1-(4-methoxy-phenyl)-ethanone (13.8 g, 36.75 mmol, see Preparation 62) in MeOH (150 ml) cooled to  $0^\circ\text{C}$  was added  $\text{NaBH}_4$  in five equal portions over 30 minutes. The reaction was stirred overnight while slowly warming to RT. Water (100 ml) was added to the reaction mixture, the volume was reduced by half *in vacuo*, and the mixture extracted with  $\text{CH}_2\text{Cl}_2$  (3x100 ml). The combined organics were dried over  $\text{MgSO}_4$ , filtered, and concentrated *in vacuo*. Flash chromatography of the residue ( $\text{SiO}_2$ , gradient elution 20%-50%  $\text{EtOAc}/\text{hexanes}$ ) gave the title product (12.87 g, 34.1 mmol, 93% yield).

**[0207]** Alternatively, to a solution of 2-benzylamino-1-(4-methoxy-phenyl)-ethanol (1.20 g, 4.66 mmol, see Preparation 61) and 3-methoxybenzaldehyde (0.700 g, 5.13 mmol) in 1,2 DCE (25 ml) was added  $\text{NaB}(\text{OAc})_3\text{H}$ . The mixture was stirred at RT overnight. The reaction mixture was poured into sat.  $\text{NaHCO}_3$  (30 ml) and extracted with  $\text{CH}_2\text{Cl}_2$  (3x50 ml). The combined organics were dried over  $\text{MgSO}_4$ , filtered, and concentrated *in vacuo*. The residue was purified via flash chromatography (Biotage 15%  $\text{EtOAc}/\text{hexanes}$ ) to give the title product (1.72 g, 4.56 mmol, 98% yield).

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  2.58 (1H, m), 3.44 (1H, m), 3.72 (1H, m), 3.75 (3H, s), 3.78 (3H, s), 3.88 (1H, m), 4.65 (1H, m), 6.79-6.90 (5H, m) and 7.12-7.32 (8H, m). MS 378 (M+1).

**Preparation 64****7-Methoxy-4-(4-methoxy-phenyl)-1,2,3,4-tetrahydroisoquinoline**

[0208] A suspension of 10% Pd/C (3.80 g) and 2-benzyl-7-methoxy-4-(4-methoxyphenyl)-1,2,3,4-tetrahydroisoquinoline (3.87 g, 10.8 mmol, see Example 80) in EtOH (160 ml) was hydrogenated at 50 p.s.i. for 12 hours. The catalyst was removed via filtration through diatomaceous earth and the filtrate was concentrated *in vacuo*.

Purification via flash chromatography (Biotage, SiO<sub>2</sub>, 9:1 EtOAc: MeOH then 1:1 EtOAc: MeOH) gave the title product (2.19 g, 8.13 mmol, 76% yield).

<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ<sub>H</sub> 2.88 (1H, dd, J 12.5 and 8.5 Hz), 3.21-3.31 (2H, m), 3.71 (3H, s), 3.73 (3H, s), 3.91-4.06 (2H, m), 6.61 (2H, m), 6.68 (1H, m), 6.81 (2H, d J 8.5 Hz) and 6.98 (2H, d, J 8.5 Hz). MS 270 (M+1).

**Preparation 65****6-Methoxy-1-(4-methoxy-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinoline**

[0209] A stirred solution of 4-methoxy-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide in POCl<sub>3</sub> under an atmosphere of N<sub>2</sub> was heated at reflux for 14 hours. The solution was allowed to cool to rt and concentrated in *vacuo*.

The residue was taken up in CH<sub>2</sub>Cl<sub>2</sub> (30 ml) and washed with sat. NaHCO<sub>3</sub> until CO<sub>2</sub> evolution ceased. The organic layer was dried over MgSO<sub>4</sub>, filtered, and concentrated in *vacuo*. The residue was dissolved in MeOH (20 ml), cooled to 0°C, and NaBH<sub>4</sub> (0.105g, 2.77 mmol) was added in a portionwise manner. The reaction was stirred for 2 hours at 0°C, then additional NaBH<sub>4</sub> (0.105g, 2.77 mmol) was added. The reaction was stirred at rt overnight, diluted with CH<sub>2</sub>Cl<sub>2</sub> (50 ml), and washed with sat. NaHCO<sub>3</sub> solution (1x 50 ml). The organic layer was dried over MgSO<sub>4</sub>, filtered, and concentrated in *vacuo*. The residue was subjected to flash chromatography (SiO<sub>2</sub>, hexanes:EtOAc 10:1) to give the desired compound as a colorless oil (0.550g, 1.59 mmol, 58% yield).

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 2.89 (2H, m), 3.46 (1H, m), 3.63 (1H, m), 3.73 (3H, s), 3.77 (3H, s), 5.73 (1H, s) 6.69-7.22 (12H, overlapping m).

[0210] See also Nagarajan et al., *Ind. J. Chem.*, 24B:83-97 (1985).

**Preparation 66****6-Methoxy-1,2-diphenyl-1,2,3,4-tetrahydroisoquinoline**

[0211] The title compound was prepared by analogy to Preparation 65 except that N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide was used instead of 4-methoxy-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide and the final residue was subjected to flash column chromatography using hexanes:EtOAc 10:1.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 2.88 (2H, m), 3.47 (1H, m), 3.69 (1H, m), 3.77 (3H, s), 5.77 (1H, s), 6.70-7.24 (13H, overlapping m). MS 316 (M+1).

[0212] See also Nagarajan et al., *Ind. J. Chem.*, 24B:83-97 (1985).

**Preparation 67****1-(4-Chloro-phenyl)-6-methoxy-2-phenyl-1,2,3,4-tetrahydroisoquinoline**

[0213] The title compound was prepared by analogy to Preparation 65 except that 4-chloro-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide was used instead of 4-methoxy-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide, and the final residue was not subjected to flash chromatography. MS 349 (M+1).

[0214] See also Nagarajan et al., *Ind. J. Chem.*, 24B:83-97 (1985).

**Example 1****2,2,2-Trifluoro-1-[1-(4-hydroxyphenyl)-6-methoxy-3,4-dihydro-1H-isoquinolin-2-yl]ethanone**

[0215] 1-[1-(4-Benzyloxyphenyl)-6-methoxy-3,4-dihydro-1H-isoquinolin-2-yl]-2,2,2-trifluoroethanone (26.97 g, 61.1 mmol) was hydrogenated using 10% Pd/C (1.000 g) in EtOH (300 ml) at an initial pressure of 44 psi (~300,000 Pa) at rt for 6 hr. The reaction mixture was filtered through a pad of Celite to remove the catalyst and the filtrate was evaporated *in vacuo* to give 18.98 g (88% yield) of white solid.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.08 - 7.06 (m, 2H), 6.96 - 6.92 (m, 1H), 6.79 - 6.69 (m, 5H), 4.81 (br s, 1H), 3.92 - 3.88

(m, 1H), 3.80 (s, 3H), 3.48 - 3.40 (m, 1H), 3.10 - 3.01 (m, 1H), 2.85 - 2.81 (m, 1H); MS m/e 352 (M<sup>+</sup>+1).

### Example 2

#### 2,2,2-Trifluoro-1-[6-methoxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinolin-2-yl]ethanone

[0216] A solution of 2,2,2-trifluoro-1-[1-(4-hydroxyphenyl)-6-methoxy-3,4-dihydro-1H-isoquinolin-2-yl]ethanone (18.98 g, 54.0 mmol) in anhydrous DMF (200 ml) was added to a suspension of NaH (2.592 g, 108.0 mmol) in anhydrous DMF (400 ml) at rt under N<sub>2</sub>. After stirring at rt for 1 hr, a solution of 1-(2-chloroethyl)pyrrolidine hydrochloride (9.184 g, 54.0 mmol) in anhydrous DMF (200 ml) was added and the reaction mixture was heated to 100°C for 4 hr. The reaction mixture was cooled to rt, diluted with H<sub>2</sub>O (2000 ml), and extracted with EtOAc (4 x 250 ml). The combined extracts were washed with H<sub>2</sub>O (3 x 250 ml), dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to give 21.15 g of yellow oil. Purification by flash chromatography, eluting with EtOAc: MeOH (8:2) gave 10.503 g (43% yield) of yellow oil.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.11 - 7.09 (m, 2H), 6.93 (d, J = 8.51 Hz, 1 H), 6.81 - 6.79 (m, 2H), 6.77 - 6.73 (m, 1H), 6.71 - 6.69 (m, 2H), 4.16 (br s, 2H), 3.91 - 3.87 (m, 1H), 3.79 (s, 3H), 3.47 - 3.40 (m, 1H), 3.10 - 2.94 (m, 3H), 2.86 - 2.77 (m, 5H), 1.86 (s, 4H); MS m/e 449 (M<sup>+</sup>+1).

### Example 3

#### 2,2,2-Trifluoro-1-[6-hydroxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinolin-2-yl]ethanone

[0217] A 1M solution of BBr<sub>3</sub> in CH<sub>2</sub>Cl<sub>2</sub> (35.0 ml, 35.0 mmol) was added slowly to a solution of 2,2,2-trifluoro-1-[6-methoxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinolin-2-yl]ethanone (6.26 g, 14.0 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (400 ml) at 0°C under N<sub>2</sub>. The reaction mixture was warmed to rt and stirred at rt for 21 hr.

MeOH (100 ml) was slowly added with stirring and the resulting solution was concentrated *in vacuo* to a red oil. Purification by flash chromatography, eluting with EtOAc: MeOH (8:2), gave 1.886 g (31% yield) of yellow solid.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.07 (d, J = 8.30 Hz, 2H), 6.79 (d, J = 8.30 Hz, 1H), 6.71 - 6.63 (m, 5H), 4.24 (br s, 2H), 3.85 - 3.81 (m, 1 H), 3.37 - 3.31 (m, 1 H), 3.20 - 2.95 (m, 7H), 2.71 - 2.66 (m, 1 H), 1.98 (s, 4H); MS m/e 435 (M<sup>+</sup>+1).

### Example 4

#### {6-Hydroxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinolin-2-yl}-phenylmethanone

[0218] A mixture of {6-benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-3,4-dihydro-1H-isoquinolin-2-yl}phenylmethanone (0.043 g, 0.08 mmol), ammonium formate (0.127 g, 2.01 mmol), and 20% Pd(OH)<sub>2</sub>/C (0.025 g) in MeOH (10 ml) was refluxed under N<sub>2</sub> for 1 hr, then filtered through Celite. The filtrate was evaporated *in vacuo* to a white residue that was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (10 ml) and washed with 1M NaOH (5 ml). The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 10 ml) and all of the combined organic solutions were dried over MgSO<sub>4</sub> and concentrated *in vacuo* to give 0.028 g (78% yield) of colorless residue.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.38 - 7.34 (m, 6H), 7.18 (d, J = 7.48 Hz), 6.91 - 6.62 (m, 6H), 4.24 - 4.22 (m, 2H), 3.72 - 3.52 (m, 1H), 3.26 - 2.89 (m, 8H), 2.56 - 2.52 (m, 1H), 1.95 (br s, 4H); MS m/e 443 (M<sup>+</sup>+1).

### Example 5

#### 1-[6-Hydroxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinolin-2-yl]-2,2-dimethylpropan-1-one

[0219] A mixture of 1-[6-benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinolin-2-yl]-2,2-dimethylpropan-1-one, ammonium formate (0.033 g, 0.52 mmol), and 20% Pd(OH)<sub>2</sub>/C (0.002 g) in MeOH (10 ml) was refluxed under N<sub>2</sub> for 1.5 hr, then filtered to remove the catalyst. The filtrate was evaporated *in vacuo* to a residue that was suspended in H<sub>2</sub>O (0.4 ml) and saturated NaHCO<sub>3</sub> (0.4 ml). The aqueous mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 0.8 ml) and all of the combined organic solutions were evaporated to give the crude product. Purification by reverse-phase HPLC on a Primesphere C-18HC (50.0 mm x 10.0 mm column with 5 μm particle size) column, eluting with a linear gradient starting at time 0 min. of H<sub>2</sub>O: CH<sub>3</sub>CN: 1%TFA/H<sub>2</sub>O (85: 10: 5), increasing to H<sub>2</sub>O: CH<sub>3</sub>CN: 1%TFA/H<sub>2</sub>O (5: 90: 5) at 8 min., detected on a Micromass Platform 2 mass spectrometer (DAD 190 - 600 nM) gave material which, after evaporation to dryness, was purified by reverse-phase HPLC on a Primesphere C-18HC (3.0 mm x 2.0 mm column with 5 μm particle size) column, eluting with a linear gradient starting at time 0 min. of H<sub>2</sub>O: CH<sub>3</sub>CN: TFA (99.9: 0: 0.1), increasing to H<sub>2</sub>O: CH<sub>3</sub>CN: TFA (0: 99.9: 0.1) at 4 min., detected with a UV detector (300 nM +/- 90

nM and 254 nM +/- 25 nM), gave an eluent that was evaporated to dryness to give the pure product.  
MS m/e 423 (M<sup>+</sup>+1).

#### Example 6

##### Cyclohexyl-{6-hydroxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinolin-2-yl}methanone

[0220] The title compound was prepared in a two-step sequence. The first step was performed by analogy to Preparation 8, except that trimethylacetyl chloride was replaced with cyclohexanecarbonyl chloride.

MS m/e 539 (M<sup>+</sup>+1).

[0221] The second step was performed by analogy to Example 5.

MS m/e 449 (M<sup>+</sup>+1).

#### Example 7

##### 1-{6-Hydroxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinolin-2-yl}-3-phenyl-propan-1-one

[0222] The title compound was prepared in a two-step sequence. The first step was performed by analogy to Preparation 8, except that trimethylacetyl chloride was replaced with cinnamoyl chloride.

MS m/e 559 (M<sup>+</sup>+1).

[0223] The second step was performed by analogy to Example 5.

MS m/e 471 (M<sup>+</sup>+1).

#### Example 8

##### 1-{6-Hydroxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinolin-2-yl}octan-1-one

[0224] The title compound was prepared in a two-step sequence. The first step was performed by analogy to Preparation 8, except that trimethylacetyl chloride was replaced with octanoyl chloride.

MS m/e 555 (M<sup>+</sup>+1).

[0225] The second step was performed by analogy to Example 5.

MS m/e 465 (M<sup>+</sup>+1).

#### Example 9

##### {6-Hydroxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinolin-2-yl}naphthalen-1-yl-methanone

[0226] The title compound was prepared in a two-step sequence. The first step was performed by analogy to Preparation 8, except that trimethylacetyl chloride was replaced with 1-naphthoyl chloride.

MS m/e 583 (M<sup>+</sup>+1).

[0227] The second step was performed by analogy to Example 5.

MS m/e 493 (M<sup>+</sup>+1).

#### Example 10

##### {6-Hydroxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinolin-2-yl}-(3-methoxyphenyl)methanone

[0228] The title compound was prepared in a two-step sequence. The first step was performed by analogy to Preparation 8, except that trimethylacetyl chloride was replaced with 3-methoxybenzoyl chloride.

MS m/e 563 (M<sup>+</sup>+1).

[0229] The second step was performed by analogy to Example 5.

MS m/e 473 (M<sup>+</sup>+1).

**Example 11****3-Cyclopentyl-1-[6-hydroxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinolin-2-yl]propan-1-one**

**[0230]** The title compound was prepared in a two-step sequence. The first step was performed by analogy to Preparation 8, except that trimethylacetyl chloride was replaced with 3-cyclopentylpropionyl chloride.

MS m/e 553 ( $M^{+}+1$ ).

**[0231]** The second step was performed by analogy to Example 5.

MS m/e 463 ( $M^{+}+1$ ).

**Example 12****1-[6-Hydroxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-3,4-dihydro-1H-isoquinolin-2-yl]-2,2-diphenyl-ethanone**

**[0232]** The title compound was prepared in a two-step sequence. The first step was performed by analogy to Preparation 8, except that trimethylacetyl chloride was replaced with diphenylacetyl chloride.

MS m/e 533 ( $M^{+}+1$ ).

**[0233]** The second step was performed by analogy to Example 5.

MS m/e 533 ( $M^{+}+1$ ).

**Example 13****2,2,2-Trifluoro-1-[6-hydroxy-1-(4-hydroxyphenyl)-3,4-dihydro-1H-isoquinolin-2-yl]ethanone**

**[0234]** A 1M solution of  $\text{BBr}_3$  in  $\text{CH}_2\text{Cl}_2$  (6.0 ml, 6.0 mmol) was added slowly to a solution of 1-[1-(4-benzyloxyphenyl)-6-methoxy-3,4-dihydro-1H-isoquinolin-2-yl]-2,2,2-trifluoroethanone (1.054 g, 2.4 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (100 ml) at  $0^\circ\text{C}$  under  $\text{N}_2$ . The reaction mixture was warmed to rt and stirred at rt for 17 hr. MeOH (50 ml) was slowly added with stirring and the resulting solution was concentrated *in vacuo* to a red oil. This was dissolved in MeOH (20 ml) and 1M HCl (20 ml), and the resulting solution was stirred at rt for 4 hr, then neutralized with saturated  $\text{NaHCO}_3$  solution (60 ml). The resulting suspension was concentrated *in vacuo* and the remaining material was extracted with EtOAc (4 x 30 ml). The combined extracts were dried over  $\text{MgSO}_4$  and evaporated *in vacuo* to give 0.775 g of crude brown solid. Purification by flash chromatography, eluting with hexane : EtOAc (1:1) gave 0.080 g (10% yield) of white solid.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.06 (d, J = 8.72 Hz, 2H), 6.88 (d, J = 8.30 Hz, 1H), 6.74 - 6.66 (m, 5H), 3.84 - 3.81 (m, 1H), 3.45 - 3.40 (m, 1H), 3.02 - 2.94 (m, 1H), 2.77-2.73 (m, 1H); MS m/e 336 ( $M^{+}-1$ ).

**Example 14****2,2,2-Trifluoro-1-[6-hydroxy-1-(4-hydroxyphenyl)-3,4-dihydro-1H-isoquinolin-2-yl]-ethanone**

**[0235]** The racemic material obtained in Example 13 (0.72 g) was separated on a Chiralpak AD $\text{C}$  column (Chiral Technologies, Inc., Exton, PA), eluting with hexane : isopropanol (9:1) to give the two separate enantiomers. The enantiomer with a retention time of 14.1 minutes was collected and evaporated *in vacuo* to give 0.018 g (25% recovery) of white solid.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.06 (d, J = 8.72 Hz, 2H), 6.88 (d, J = 8.30 Hz, 1H), 6.74 - 6.66 (m, 5H), 3.84 - 3.81 (m, 1H), 3.45 - 3.40 (m, 1H), 3.02 - 2.94 (m, 1H), 2.77 - 2.73 (m, 1H); MS m/e 336 ( $M^{+}-1$ ).

**Example 15****2,2,2-Trifluoro-1-[1-(4-hydroxyphenyl)-3,4-dihydro-1H-isoquinolin-2-yl]ethanone**

**[0236]** A mixture of 1-[1-(4-benzyloxyphenyl)-3,4-dihydro-1H-isoquinolin-2-yl]-2,2,2-trifluoroethanone (0.144 g, 0.35 mmol), ammonium formate (0.441 g, 7.00 mmol), and 20%  $\text{Pd}(\text{OH})_2/\text{C}$  (0.100 g) in MeOH (50 ml) was refluxed under  $\text{N}_2$  for 1.5 hr, then filtered through Celite. The filtrate was evaporated *in vacuo* to a yellow residue that was suspended in a mixture of  $\text{H}_2\text{O}$  (15 ml) and saturated  $\text{NaHCO}_3$  solution (15 ml), then extracted with  $\text{CHCl}_3$  (4 x 15 ml). The combined organic extracts were dried over  $\text{MgSO}_4$  and concentrated *in vacuo* to give 0.115 g of colorless oil. Purification by flash chromatography, eluting with hexane : EtOAc (8:2) gave 0.100 g (89% yield) of colorless oil.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.29 - 7.16 (m, 4H), 7.09 - 7.00 (m, 3H), 6.77 - 6.72 (m, 2H), 3.96 - 3.91 (m, 1H), 3.52 -

3.44 (m, 1H), 3.14 - 3.05 (m, 1H), 2.90 - 2.86 (m, 1H); MS m/e 320 ( $M^+ - 1$ ).

#### Example 16

##### 2,2,2-Trifluoro-1-[1-(4-hydroxyphenyl)-3,4-dihydro-1H-isoquinolin-2-yl]ethanone

[0237] The racemic material obtained in Example 15 (0.80 g) was separated on a Chiralpak AD column, eluting with hexane : isopropanol (9:1) to give the two separate enantiomers. The enantiomer with a retention time of 9.7 minutes was collected and evaporated *in vacuo* to give 0.030 g (38% recovery) of white solid.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.29 - 7.16 (m, 4H), 7.09 - 7.00 (m, 3H), 6.77 - 6.72 (m, 2H), 3.95 - 3.91. (m, 1H), 3.51 - 3.43 (m, 1H), 3.14 - 3.05 (m, 1H), 2.90 - 2.85 (m, 1H); MS m/e 320 ( $M^+ - 1$ ).

#### Example 17

##### 2,2,2-Trifluoro-1-[1-(4-hydroxyphenyl)-6-(2-pyrrolidin-1-yl-ethoxy)-3,4-dihydro-1H-isoquinolin-2-yl]ethanone

[0238] A solution of 1M TBAF in THF (2.8 ml, 2.82 mmol) and toluene-4-sulfonic acid 4-[6-(2-pyrrolidin-1-yl-ethoxy)-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinolin-1-yl]phenyl ester (0.415 g, 0.71 mmol) in anhydrous THF (50 ml) was refluxed under N<sub>2</sub> for 18 hr, then concentrated *in vacuo* to a green oil. This was suspended in sat. NaHCO<sub>3</sub> solution (25 ml), then extracted with EtOAc (3 x 40 ml). The combined extracts were dried over MgSO<sub>4</sub> and concentrated *in vacuo* to give 0.566 g of green oil. Purification by flash chromatography, eluting with EtOAc : MeOH (8:2) gave 0.241 g (79% yield) of yellow oil.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.06 - 6.99 (m, 2H), 6.94 - 6.87 (m, 1H), 6.83 (d, J = 8.51 Hz, 1H), 6.71 (d, J = 8.51 Hz, 1H), 6.66 (m, 1H), 6.64 - 6.57 (m, 2H), 4.23 - 4.21 (m, 2H), 3.90 - 3.85 (m, 1H), 3.46 - 3.35 (m, 1H), 3.13 (s, 2H), 3.05 - 2.95 (m, 5H), 2.79-2.74 (m, 1H), 1.95 (s, 4H); MS m/e 435 ( $M^+ + 1$ ).

#### Example 18

##### 2,2,2-Trifluoro-1-[1-(4-iodophenyl)-6-methoxy-3,4-dihydro-1H-isoquinolin-2-yl]ethanone

[0239] Trifluoroacetic anhydride (1.76 g, 8.36 mmol) was added to a solution of 1-(4-iodophenyl)-6-methoxy-1,2,3,4-tetrahydroisoquinoline (2.35 g, 6.44 mmol) and Et<sub>3</sub>N (1.0 ml, 7.1 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (50 ml) at 0°C under N<sub>2</sub>. The resulting yellow solution was stirred at 0°C for 2 hr, then washed first with 1 M HCl (2 x 25 ml) followed by 50% saturated NaHCO<sub>3</sub> solution (2 x 25 ml), dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated *in vacuo* to give a yellow oil. Purification by flash chromatography, eluting with hexane : EtOAc (6:1) gave 1.87 g (63% yield) of yellow oil.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.62 (d, J = 8.31 Hz, 2H), 6.97 - 6.91 (m, 3H), 6.79 - 6.76 (m, 1H), 6.71 - 6.69 (m, 2H), 3.95 - 3.91 (m, 1H), 3.81 (s, 3H), 3.45 - 3.37 (m, 1H), 3.07 - 3.02 (m, 1 H), 2.87 - 2.81 (m, 1 H) MS m/e 460 ( $M^+ - 1$ ).

#### Example 19

##### 2,2,2-Trifluoro-1-[6-hydroxy-1-(4-iodophenyl)-3,4-dihydro-1H-isoquinolin-2-yl]ethanone

[0240] A 1M solution of BBr<sub>3</sub> in CH<sub>2</sub>Cl<sub>2</sub> (0.49 ml, 0.49 mmol) was added slowly to a solution of 2,2,2-trifluoro-1-[1-(4-iodophenyl)-6-methoxy-3,4-dihydro-1 H-isoquinolin-2-yl]ethanone (0.150 g, 0.33 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (10 ml) at 0°C under N<sub>2</sub>. The reaction mixture was stirred at 0°C for 1 hr, then warmed to rt and stirred at rt for 19 hr. MeOH (10 ml) was slowly added with stirring and the resulting solution was concentrated *in vacuo* to a yellow oil. Purification by flash chromatography, eluting with hexane : EtOAc (8:2) gave 0.121 g (83% yield) of yellow solid.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.62 (d, J = 8.51 Hz, 2H), 6.94 (d, J = 8.51 Hz, 2H), 6.87 (d, J = 8.30 Hz, 1H), 6.71 - 6.68 (m, 3H), 3.94 - 3.89 (m, 1H), 3.44 - 3.36 (m, 1H), 3.08 - 2.99 (m, 1 H), 2.83 - 2.78 (m, 1 H), MS m/e 446 ( $M^+ - 1$ ).

#### Example 20

##### 1-(1-Cyclohexyl-6-methoxy-3,4-dihydro-1H-isoquinolin-2-yl)-2,2,2-trifluoroethanone

[0241] Trifluoroacetic anhydride (0.128 g, 0.61 mmol) was added to a solution of 1-cyclohexyl-6-methoxy-1,2,3,4-tetrahydroisoquinoline (0.150 g, 0.61 mmol) and Et<sub>3</sub>N (0.17 ml, 0.122 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (20 ml) at 0°C under N<sub>2</sub>. The resulting yellow solution was stirred at rt for 22 hr, washed first with 1M HCl (5 ml) then 1M NaHCO<sub>3</sub> (5 ml), dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to give 0.248 g of yellow oil. Purification by flash chromatography, eluting



with hexane : EtOAc (19:1) gave 0.198 g (95% yield) of yellow oil.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.02 (d, J = 8.51 Hz, 1H), 6.72 (dd, J = 8.51 Hz, J = 2.70 Hz, 1H), 6.65 (d, J = 2.49 Hz, 1H), 5.16 (d, J = 9.13 Hz, 1H), 3.97 - 3.94 (m, 1H), 3.77 (s, 3H), 3.75 - 3.68 (m, 1H), 3.09 - 2.88 (m, 2H), 1.75 (br s, 4H), 1.11 - 1.02 (m, 5H); MS m/e 342 (M<sup>+</sup>+1).

## Example 21

### 1-(1-Cyclohexyl-6-hydroxy-3,4-dihydro-1H-isoquinolin-2-yl)-2,2,2-trifluoroethanone

[0242] A 1 M solution of BBr<sub>3</sub> in CH<sub>2</sub>Cl<sub>2</sub> (0.77 ml, 0.77 mmol) was added slowly to a solution of 1-(1-cyclohexyl-6-methoxy-3,4-dihydro-1H-isoquinolin-2-yl)-2,2,2-trifluoroethanone (0.176 g, 0.52 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (10 ml) at 0°C under N<sub>2</sub>.

The reaction mixture was stirred at 0°C for 30 min, then warmed to rt and stirred at rt for 21 hr. MeOH (10 ml) was slowly added with stirring and the resulting solution was concentrated *in vacuo* to a orange oil. Purification by flash chromatography, eluting with hexane : EtOAc (8:2) gave 0.193 g (100% yield) of white solid.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 6.97 (d, J = 8.52 Hz, 1H), 6.66 - 6.60 (m, 3H), 5.15 (d, J = 8.93 Hz, 1H), 4.69 (br s, 1H), 3.97 - 3.94 (m, 1H), 3.75 - 3.67 (m, 1H), 3.04 - 2.87 (m, 3H), 1.75 (br s, 4H), 1.12 - 1.10 (m, 5H); MS m/e 326 (M<sup>+</sup>-1).

## Example 22

### 2,2,2-Trifluoro-1-(6-methoxy-1-phenyl-3,4-dihydro-1H-isoquinolin-2-yl)ethanone

[0243] Trifluoroacetic anhydride (0.089 ml, 0.627 mmol) was added to a solution of 6-methoxy-1-phenyl-1,2,3,4-tetrahydroisoquinoline (0.150 g, 0.627 mmol) and Et<sub>3</sub>N (0.174 ml, 1.25 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (20 ml) at 0°C under N<sub>2</sub>. The resulting solution was warmed to rt and stirred at rt for 68 hr, washed with 1 M HCl (5 ml), then with sat. NaHCO<sub>3</sub> solution (5 ml), dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to give 0.216 g of colorless oil. Purification by flash chromatography, eluting with hexane : EtOAc (19:1) gave 0.188 g (89% yield) of colorless oil.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.30 - 7.28 (m, 3H), 7.21 - 7.18 (m, 2H), 6.96 (d, J = 8.72 Hz, 1H), 6.79 - 6.76 (m, 2H), 6.72 (s, 1H), 3.93 - 3.90 (m, 1H), 3.81 (s, 3H), 3.50 - 3.42 (m, 1H), 3.08 - 3.03 (m, 1H), 2.86 - 2.82 (m, 1H); MS m/e 334 (M<sup>+</sup>-1).

## Example 23

### 2,2,2-Trifluoro-1-(6-hydroxy-1-phenyl-3,4-dihydro-1H-isoquinolin-2-yl)ethanone

[0244] A 1M solution of BBr<sub>3</sub> in CH<sub>2</sub>Cl<sub>2</sub> (0.77 ml, 0.77 mmol) was added slowly to a solution of 2,2,2-trifluoro-1-(6-methoxy-1-phenyl-3,4-dihydro-1H-isoquinoline-2-yl)ethanone (0.172 g, 0.513 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (10 ml) at 0°C under N<sub>2</sub>. The resulting yellow solution was stirred at 0°C for 30 min, then warmed to rt and stirred at rt for 22 hr. MeOH (5 ml) was slowly added to the reaction mixture with stirring and the resulting solution was concentrated *in vacuo* to give 0.242 g of yellow oil.

Purification by flash chromatography, eluting with hexane : EtOAc (8:2) gave 0.143 g (87% yield) of white solid.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.30 - 7.27 (m, 3H), 7.20 - 7.17 (m, 2H), 6.91 (d, J = 8.09 Hz, 1H), 6.76 (s, 1H), 6.70 - 6.74 (m, 2H), 4.76 (br s, 1H), 3.92 - 3.88 (m, 1H), 3.49 - 3.41 (m, 1H), 3.09 - 2.97 (m, 1H), 2.83 - 2.79 (m, 1H); MS m/e 320 (M<sup>+</sup>-1).

## Example 24

### 2,2,2-Trifluoro-1-(6-methoxy-1-thiophen-2-yl-3,4-dihydro-1H-isoquinolin-2-yl)ethanone

[0245] Trifluoroacetic anhydride (0.70 ml, 4.97 mmol) was added to a solution of 6-methoxy-1-thiophen-2-yl-1,2,3,4-tetrahydroisoquinoline (1.220 g, 4.97 mmol) and Et<sub>3</sub>N (1.39 ml, 9.94 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (40 ml) at 0°C under N<sub>2</sub>. The resulting solution was warmed to rt and stirred at rt for 68 hr, washed with 1M HCl (20 ml) followed by sat. NaHCO<sub>3</sub> solution (20 ml), dried over MgSO<sub>4</sub>, then concentrated *in vacuo* to give 1.863 g of red oil. Purification by flash chromatography, eluting with hexane : EtOAc (19:1) gave 1.340 g (79% yield) of yellow oil.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.24 - 7.23 (m, 1H), 7.09 (d, J = 8.72 Hz, 1H), 6.92 - 6.90 (m, 1H), 6.84 - 6.83 (m, 2H), 6.83 - 6.77 (m, 1H), 6.69 (s, 1H), 3.98 - 3.95 (m, 1H), 3.81 (s, 3H), 3.62 - 3.55 (m, 1H), 3.11 - 3.02 (m, 1H), 2.88 - 2.84 (m, 1H); MS m/e 340 (M<sup>+</sup>-1).

**Example 25****2,2,2-Trifluoro-1-(6-hydroxy-1-thiophen-2-yl-3,4-dihydro-1H-isoquinolin-2-yl)ethanone**

- 5 **[0246]** A 1M solution of  $\text{BBr}_3$  in  $\text{CH}_2\text{Cl}_2$  (0.66 ml, 0.66 mmol) was added slowly to a solution of 2,2,2-trifluoro-1-(6-methoxy-1-thiophen-2-yl-3,4-dihydro-1H-isoquinolin-2-yl)ethanone (0.150 g, 0.439 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (10 ml) at  $0^\circ\text{C}$  under  $\text{N}_2$ . The resulting red solution was stirred at  $0^\circ\text{C}$  for 30 min, then warmed to rt and stirred at rt for 20 hr. MeOH (5 ml) was slowly added to the reaction mixture with stirring and the resulting solution was concentrated *in vacuo* to give 0.138 g of a black residue.
- 10 Purification by flash chromatography, eluting with hexane : EtOAc (8:2) gave 0.006 g (4% yield) of white solid.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.24 (s, 1H), 7.04 (d,  $J$  = 8.72 Hz, 1H), 6.90 - 6.89 (m, 1H), 6.82 (m, 2H), 6.69 (d,  $J$  = 8.92 Hz, 1H), 6.65 (s, 1H), 3.96 - 3.93 (m, 1H), 3.60 - 3.54 (m, 1H), 3.03 - 3.00 (m, 1H), 2.84 - 2.80 (m, 1H); MS  $m/e$  326 ( $\text{M}^+ - 1$ ).

**Example 26****1-[1-(4-Bromophenyl)-6-methoxy-3,4-dihydro-1H-isoquinolin-2-yl]-2,2,2-trifluoroethanone**

- [0247]** Trifluoroacetic anhydride (0.067 ml, 0.471 mmol) was added to a solution of 1-(4-bromophenyl)-6-methoxy-1,2,3,4-tetrahydroisoquinoline (0.150 g, 0.471 mmol) and  $\text{Et}_3\text{N}$  (0.13 ml, 0.942 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (20 ml) at  $0^\circ\text{C}$  under  $\text{N}_2$ . The resulting solution was warmed to rt and stirred at rt for 68 hr, washed with 1 M HCl (5 ml) and then with sat.  $\text{NaHCO}_3$  solution (5 ml), dried over  $\text{MgSO}_4$ , and concentrated *in vacuo* to give 0.220 g of yellow oil. Purification by flash chromatography, eluting with hexane : EtOAc (19:1) gave 0.181 g (93% yield) of white solid.
- 25  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.41 (d,  $J$  = 8.51 Hz, 2H), 7.07 (d,  $J$  = 8.51 Hz, 2H), 6.91 (d,  $J$  = 8.51 Hz, 1H), 6.77 (dd,  $J$  = 8.30 Hz,  $J$  = 2.49 Hz, 1H), 6.70 (s, 2H), 3.93 - 3.81 (m, 1H), 3.80 (s, 3H), 3.44 - 3.67 (m, 1H), 3.11-3.02 (m, 1H), 2.85-2.82 (m, 1H); MS  $m/e$  412, 414 bromine isotope pattern ( $\text{M}^+ - 1$ ).

**Example 27****1-[1-(4-Bromophenyl)-6-hydroxy-3,4-dihydro-1H-isoquinolin-2-yl]-2,2,2-trifluoroethanone**

- [0248]** A 1M solution of  $\text{BBr}_3$  in  $\text{CH}_2\text{Cl}_2$  (0.61 ml, 0.61 mmol) was added slowly to a solution of 1-[1-(4-bromophenyl)-6-methoxy-3,4-dihydro-1H-isoquinolin-2-yl]-2,2,2-trifluoroethanone (0.168 g, 0.406 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (10 ml) at  $0^\circ\text{C}$  under  $\text{N}_2$ .
- 35 The resulting orange solution was stirred at  $0^\circ\text{C}$  for 30 min, then warmed to rt and stirred at rt for 21 hr. MeOH (5 ml) was slowly added to the reaction mixture with stirring and the resulting solution was concentrated *in vacuo* to give 0.201 g of yellow residue. Purification by flash chromatography, eluting with hexane : EtOAc (8:2) gave 0.149 g (92% yield) of white solid.
- 40  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.44 - 7.41 (m, 2H), 7.09 - 7.07 (m, 2H), 6.88 (d,  $J$  = 8.10 Hz, 1H), 6.71 - 6.68 (m, 3H), 3.94 - 3.90 (m, 1H), 3.43 - 3.36 (m, 1H), 3.09 - 2.98 (m, 1H), 2.83 - 2.79 (m, 1H); MS  $m/e$  400, 398 bromine isotope pattern ( $\text{M}^+ - 1$ ).

**Example 28****2-Benzenesulfonyl-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol**

- [0249]** A mixture of 2-benzenesulfonyl-6-benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinoline (0.043 g, 0.08 mmol), ammonium formate (0.096 g, 1.52 mmol), and 20%  $\text{Pd}(\text{OH})_2/\text{C}$  (0.025 g) in MeOH (10 ml) was refluxed under  $\text{N}_2$  for 1 hr, then filtered through Celite. The filtrate was evaporated *in vacuo* to a white residue that was dissolved in  $\text{CH}_2\text{Cl}_2$  (10 ml) and washed with 1M NaOH (5 ml). The aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  (3 x 10 ml), and all of the combined organic solutions were dried over  $\text{MgSO}_4$  and concentrated *in vacuo* to give 0.026 g (72% yield) of colorless residue.
- 50  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.65 (d,  $J$  = 8.09 Hz, 4H), 7.40 - 7.38 (m, 2H), 7.29 (d,  $J$  = 7.47 Hz, 4H), 6.73 (d,  $J$  = 8.51 Hz, 2H), 6.68 (d,  $J$  = 8.09 Hz, 2H), 6.56 (d,  $J$  = 9.13 Hz, 1H), 6.37 (s, 1H), 6.09 (s, 1H), 4.21 (br s, 2H), 3.69 - 3.65 (m, 1H), 3.17 - 3.13 (m, 4H), 2.48 - 2.44 (m, 1 H), 2.35 - 2.29 (m, 1 H), 1.96 (br s, 4H); MS  $m/e$  479 ( $\text{M}^+ + 1$ ).
- 55

**Example 29****2-(Naphthalene-1-sulfonyl)-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0250]** A mixture of 6-benzyloxy-2-(naphthalene-1-sulfonyl)-1-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-1,2,3,4-tetrahydroisoquinoline (0.012 g, 0.019 mmol), ammonium formate (0.033 g, 0.52 mmol), and 20% Pd(OH)<sub>2</sub>/C (0.002 g) in MeOH (10 ml) was refluxed under N<sub>2</sub> for 1.5 hr, then filtered to remove the catalyst. The filtrate was evaporated *in vacuo* to a residue that was suspended in a mixture of H<sub>2</sub>O (0.4 ml) and sat. NaHCO<sub>3</sub> (0.4 ml). The aqueous mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 0.8 ml), and all of the combined organic solutions were evaporated to give the crude product.

Purification by reverse-phase HPLC, eluting with gradient Primesphere C-18HC (3.0 mm x 2.0 mm column with 5 μm particle size) column, eluting with a linear gradient starting at time 0 min. of H<sub>2</sub>O : CH<sub>3</sub>CN : TFA (99.9 : 0 : 0.1), increasing to H<sub>2</sub>O : CH<sub>3</sub>CN : TFA (0 : 99.9 : 0.1) at 4 min., detected with a UV detector (300 nM +/- 90 nM and 254 nM +/- 25 nM), gave an eluent that was evaporated to dryness to give the named product. MS m/e 493 (M<sup>+</sup>+1).

**Example 30****2-Phenylmethanesulfonyl-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0251]** The title compound was prepared in a two-step sequence. The first step was performed by analogy to Preparation 32, except that 1-naphthalenesulfonyl chloride was replaced with α-toluenesulfonyl chloride.

MS m/e 583 (M<sup>+</sup>+1).

**[0252]** The second step was performed by analogy to Example 29.

MS m/e 493 (M<sup>+</sup>+1).

**Example 31****2-(Butane-1-sulfonyl)-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0253]** The title compound was prepared in a two-step sequence. The first step was performed by analogy to Preparation 32, except that 1-naphthalenesulfonyl chloride was replaced with 1-butanethanesulfonyl chloride.

MS m/e 549 (M<sup>+</sup>+1).

**[0254]** The second step was performed by analogy to Example 29.

MS m/e 459 (M<sup>+</sup>+1).

**Example 32****2-Methanesulfonyl-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0255]** The title compound was prepared in a two-step sequence. The first step was performed by analogy to Preparation 32, except that 1-naphthalenesulfonyl chloride was replaced with methanesulfonyl chloride.

MS m/e 507 (M<sup>+</sup>+1).

**[0256]** The second step was performed by analogy to Example 29.

MS m/e 417 (M<sup>+</sup>+1).

**Example 33****2-(4-Propylbenzenesulfonyl)-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0257]** The title compound was prepared in a two-step sequence. The first step was performed by analogy to Preparation 32, except that 1-naphthalenesulfonyl chloride was replaced with p-n-propylbenzenesulfonyl chloride.

MS m/e 611 (M<sup>+</sup>+1).

**[0258]** The second step was performed by analogy to Example 29.

MS m/e 521 (M<sup>+</sup>+1).

**Example 34****2-(4-Isopropylbenzenesulfonyl)-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0259]** The title compound was prepared in a two-step sequence. The first step was performed by analogy to Preparation 32, except that 1-naphthalenesulfonyl chloride was replaced with 4-isopropylbenzenesulfonyl chloride.

MS m/e 611 ( $M^{+}+1$ ).

**[0260]** The second step was performed by analogy to Example 29.

MS m/e 521 ( $M^{+}+1$ ).

**Example 35****1-[4-(2-Pyrrolidin-1-yl-ethoxy)phenyl]-2-(toluene-4-sulfonyl)-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0261]** The title compound was prepared in a two-step sequence. The first step was performed by analogy to Preparation 32, except that 1-naphthalenesulfonyl chloride was replaced with 4-methylbenzenesulfonyl chloride.

MS m/e 583 ( $M^{+}+1$ ).

**[0262]** The second step was performed by analogy to Example 29.

MS m/e 493 ( $M^{+}+1$ ).

**Example 36****2-(1-Methyl-1H-imidazole-4-sulfonyl)-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0263]** The title compound was prepared in a two-step sequence. The first step was performed by analogy to Preparation 32, except that 1-naphthalenesulfonyl chloride was replaced with 1-methylimidazolesulfonyl chloride.

MS m/e 573 ( $M^{+}+1$ ).

**[0264]** The second step was performed by analogy to Example 29.

MS m/e 483 ( $M^{+}+1$ ).

**Example 37****N-(4-{6-Hydroxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinoline-2-sulfonyl}phenyl)acetamide**

**[0265]** The title compound was prepared in a two-step sequence. The first step was performed by analogy to Preparation 32, except that 1-naphthalenesulfonyl chloride was replaced with p-acetamidobenzenesulfonyl chloride.

MS m/e 626 ( $M^{+}+1$ ).

**[0266]** The second step was performed by analogy to Example 29.

MS m/e 536 ( $M^{+}+1$ ).

**Example 38****4-Amino-3-{6-hydroxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinoline-2-sulfonyl}pent-3-en-2-one**

**[0267]** A mixture of 6-benzyloxy-2-(3,5-dimethylisoxazole-4-sulfonyl)-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinoline (0.036 g, 0.061 mmol), ammonium formate (0.077 g, 1.22 mmol), and 20% Pd(OH)<sub>2</sub>/C (0.020 g) in MeOH (20 ml) was refluxed under N<sub>2</sub> for 1.5 hr, then filtered through Celite. The filtrate was evaporated *in vacuo* to a yellow residue that was suspended in a mixture of H<sub>2</sub>O (3 ml) and sat. NaHCO<sub>3</sub> solution (3 ml), then extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 x 3 ml). The combined organic extracts were dried over MgSO<sub>4</sub> and concentrated *in vacuo* to give 0.023 g (76% yield) of yellow oil. Further purification by preparative TLC, eluting with SiO<sub>2</sub>, EtOAc : MeOH (1:1), gave 0.019 g (63% yield) of colorless oil.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 11.73 - 11.65 (m, 1 H), 6.95 - 6.88 (m, 3H), 6.75 (d, J = 8.52 Hz, 1H), 6.68 (d, J = 8.10 Hz, 1H), 6.67 - 6.52 (m, 3H), 6.08 (br s, 1H), 5.80 (s, 1H), 4.05 - 4.02 (m, 2H), 3.55 - 3.52 (m, 1H), 3.10 - 2.97 (m, 2H), 2.90 - 2.84 (m, 1H), 2.79 - 2.63 (m, 4H), 2.50 (s, 1H), 2.37 (s, 3H), 2.34 (s, 3H), 1.84 - 1.79 (m, 4H); MS m/e 500 ( $M^{+}+1$ ).

**Example 39****2-(3,5-Dimethylisoxazole-4-sulfonyl)-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol.**

**[0268]** A 1 M solution of  $\text{BBr}_3$  in  $\text{CH}_2\text{Cl}_2$  (0.095 ml, 0.095 mmol) was added slowly to a solution of 6-benzyloxy-2-(3,5-dimethylisoxazole-4-sulfonyl)-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinoline (0.037 g, 0.063 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (10 ml) at  $0^\circ\text{C}$  under  $\text{N}_2$ . The reaction mixture was stirred at  $0^\circ\text{C}$  for 1 hr, then warmed to rt and stirred at rt for 4 hr. MeOH (5 ml) was slowly added with stirring and the resulting solution was concentrated *in vacuo* to an orange residue. Purification by preparative TLC, eluting with EtOAc : MeOH (1:1), and then purification again by preparative TLC, eluting with EtOAc : MeOH (8:2), gave 0.011 g (35% yield) of white solid.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  6.99 (d, J = 8.72 Hz, 2H), 6.75 (d, J = 8.51 Hz, 1H), 6.69 (d, J = 8.72 Hz, 2H), 6.61 (dd, J = 8.30 Hz, J = 2.49 Hz, 1H), 6.51 (d, J = 2.28 Hz, 1H), 6.01 (s, 1H), 4.25 (br s, 2H), 3.62 - 3.57 (m, 1H), 3.24 - 3.16 (m, 3H), 3.05 (br s, 4H), 2.66 - 2.57 (m, 1H), 2.52 (s, 3H), 2.47 - 2.42 (m, 1H), 2.18 (s, 3H), 2.00 (br s, 4H); MS m/e 498 ( $\text{M}^+ + 1$ ).

**Example 40****2-(4-Imidazol-1-yl-benzyl)-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0269]** A solution of 1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol (0.023 g, 0.067 mmol), 4-imidazol-1-yl-benzaldehyde (0.011 g, 0.063 mmol), NaOAc (0.011 g, 0.14 mmol), and sodium cyanoborohydride (0.004 g, 0.063 mmol) in MeOH (2.0 ml) was stirred at rt for 18 hr. The reaction mixture was evaporated *in vacuo* to a yellow residue that was dissolved in EtOAc (5 ml) and washed first with sat.  $\text{NaHCO}_3$  solution (5 ml) and then with  $\text{H}_2\text{O}$  (5 ml). The organic solution was dried over  $\text{MgSO}_4$  and evaporated to give 0.035 g of yellow residue. Purification by preparative TLC, eluting with EtOAc : MeOH (7:3), gave 0.007 g (23% yield) of white solid. MS m/e 495 ( $\text{M}^+ + 1$ ).

**Example 41****2-Benzo[1,3]dioxol-5-ylmethyl-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0270]** The title compound was prepared by analogy to Example 40, except that 3,4-(methylenedioxy)benzaldehyde was used in place of 4-imidazol-1-yl-benzaldehyde. MS m/e 473 ( $\text{M}^+ + 1$ ).

**Example 42****1-[4-(2-Pyrrolidin-1-yl-ethoxy)phenyl]-2-(tetrahydropyran-4-yl)-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0271]** The title compound was prepared by analogy to Example 40, except that tetrahydro-4H-pyran-4-one was used in place of 4-imidazol-1-yl-benzaldehyde.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.14 (d, J = 8.52 Hz, 2H), 6.81 - 6.79 (m, 1H), 6.75 (d, J = 8.52 Hz, 2H), 6.57 (s, 1H), 6.53 - 6.47 (m, 1H), 4.80 (s, 1H), 4.28 (br s, 2H), 4.00 - 3.97 (m, 1H), 3.91 - 3.89 (m, 1H), 3.26 - 3.18 (m, 5H), 3.11 - 3.08 (m, 2H), 2.73 - 2.65 (m, 3H), 2.00 (s, 6H), 1.80 - 1.77 (m, 3H), 1.60 - 1.57 (m, 2H), 1.43 - 1.20 (m, 1H); MS m/e 423 ( $\text{M}^+ + 1$ ).

**Example 43****2-(4,4-Dimethylcyclohexyl)-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0272]** The title compound was prepared by analogy to Example 40, except that 4,4-dimethyl-2-cyclohexene-1-one was used in place of 4-imidazol-1-yl-benzaldehyde. MS m/e 449 ( $\text{M}^+ + 1$ ).

**Example 44****2-Cyclohexyl-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0273]** The title compound was prepared by analogy to Example 40, except that cyclohexanone was used in place

of 4-imidazol-1-yl-benzaldehyde.  
MS m/e 421 ( $M^{+1}$ ).

#### Example 45

##### 2-Benzyl-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol

[0274] The title compound was prepared by analogy to Example 40, except that benzaldehyde was used in place of 4-imidazol-1-yl-benzaldehyde.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.33 - 7.21 (m, 9H), 6.76 (d,  $J$  = 8.51 Hz, 1H), 6.55 - 6.47 (m, 2H), 4.45 (s, 1H), 4.19 (br s, 2H), 3.76 (d,  $J$  = 13.70 Hz, 1H), 3.17 (d,  $J$  = 13.70 Hz, 1H), 3.04 - 2.87 (m, 9H), 2.66 (d,  $J$  = 16.39 Hz, 1H), 2.47 - 2.43 (m, 1H), 1.91 (br s, 4H); MS m/e 429 ( $M^{+1}$ ).

#### Example 46

##### 6-Hydroxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinoline-2-carboxylic acid tert-butyl ester

[0275] The title compound was prepared by analogy to Example 4, except that 6-benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-3,4-dihydro-1H-isoquinoline-2-carboxylic acid tert-butyl ester was used in place of {6-benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-3,4-dihydro-1H-isoquinolin-2-yl}phenylmethanone.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.22 - 7.06 (m, 3H), 6.81 - 6.77 (m, 1H), 6.73 (d,  $J$  = 8.72 Hz, 1H), 6.63 - 6.61 (m, 1H), 6.30 - 6.08 (m, 2H), 4.26 (br s, 2H), 3.97 - 3.93 (m, 1H), 3.17 - 2.84 (m, 8H), 2.60 - 2.56 (m, 1H), 1.98 (br s, 4H), 1.47 (s, 9H); MS m/e 439 ( $M^{+1}$ ).

#### Example 47

##### 6-Hydroxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-3,4-dihydro-1H-isoquinoline-2-carboxylic acid ethyl ester

[0276] The title compound was prepared by analogy to Example 4, except that 6-benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-3,4-dihydro-1 H-isoquinoline-2-carboxylic acid ethyl ester was used in place of {6-benzyloxy-1-[4-(2-pyrrolidin-1-yl-ethoxy)-phenyl]-3,4-dihydro-1 H-isoquinolin-2-yl}phenylmethanone.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.16 - 7.14 (m, 1 H), 7.03 (br s, 2H), 6.77 (d,  $J$  = 8.51 Hz, 1 H), 6.65 (d,  $J$  = 8.72 Hz, 2H), 6.59 - 6.57 (m, 2H), 4.14 (br s, 4H), 3.96 - 3.90 (m, 2H), 3.10 - 3.03 (m, 3H), 2.84 (br s, 5H), 2.50 (d,  $J$  = 16.19 Hz, 1H), 1.89 (br s, 4H), 1.26 (br s, 3H); MS m/e 411 ( $M^{+1}$ ).

#### Example 48

##### 1-(4-Hydroxy-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinolin-6-ol

[0277] To a stirred solution of 6-methoxy-1-(4-methoxy-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinoline (0.075g, 0.217 mmol, 1.0 eq.) in  $\text{CH}_2\text{Cl}_2$  (3 ml) cooled to  $-78^\circ\text{C}$  under an atmosphere of  $\text{N}_2$  was added  $\text{BBr}_3$  as a 1.0 M solution in  $\text{CH}_2\text{Cl}_2$  (0.65ml, 0.65 mmol, 3.0 eq.). The reaction was stirred overnight, slowly allowing the reaction mixture to warm to rt. The reaction was quenched with MeOH (2ml) and stirring was continued for 1 hour. The mixture was diluted with EtOAc (20 ml), water (2 ml), and sat.  $\text{NaHCO}_3$  (2 ml, until the pH of the aqueous layer was approximately 8). The layers were separated and the aqueous layer diluted with water (10 ml), then extracted with EtOAc (2x20ml) and  $\text{CH}_2\text{Cl}_2$  (2x20 ml). The combined organics were dried over  $\text{MgSO}_4$ , filtered, and concentrated in *vacuo*. The residue was subjected to flash chromatography ( $\text{SiO}_2$ , hexanes:EtOAc 3:2) to give the desired compound (0.053 g, 0.167 mmol, 77% yield).

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  2.85 (2H, m), 3.41 (1H, m), 3.62 (1H, m), 5.74 (1H, s), 6.62-7.14 (12H, overlapping m).

#### Example 49

##### 1-Cyclohexyl-6-methoxy-2-phenyl-1,2,3,4-tetrahydroisoquinoline

[0278] A stirred solution of cyclohexanecarboxylic acid [2-(3-methoxy-phenyl)-ethyl]-phenyl-amide in  $\text{POCl}_3$  under an atmosphere of  $\text{N}_2$  was heated at reflux for 14 hours. The solution was allowed to cool to rt and concentrated in *vacuo*. The residue was taken up in  $\text{CH}_2\text{Cl}_2$  (30 ml) and washed with sat.  $\text{NaHCO}_3$  until  $\text{CO}_2$  evolution ceased. The

organic layer was dried over  $\text{MgSO}_4$ , filtered, and concentrated in vacuo. The residue was dissolved in MeOH (20 ml), cooled to  $0^\circ\text{C}$ , and  $\text{NaBH}_4$  (0.105g, 2.77 mmol) was added in a portionwise manner. The reaction was stirred for 2 hours at  $0^\circ\text{C}$ , then additional  $\text{NaBH}_4$  (0.105g, 2.77 mmol) was added. The reaction was stirred at rt overnight, diluted with  $\text{CH}_2\text{Cl}_2$  (50 ml), and washed with sat.  $\text{NaHCO}_3$  solution (1x 50 ml). The organic layer was dried over  $\text{MgSO}_4$ , filtered, and concentrated in vacuo, to give the desired compound as a yellow oil (0.234 g, 0.729 mmol, 73% yield).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  0.84-1.18 (6H, overlapping m), 1.58-1.76 (4H, overlapping m), 1.95 (1H, m), 2.95 (2H, m), 3.43 (1H, m), 3.69 (1H, m) 3.76 (3H, s), 4.35 (1H, d, J 8.0 Hz), 6.64-7.23 (8H, overlapping m). MS 322 (M+1)

#### Example 50

##### 1-Cyclohexyl-2-phenyl-1,2,3,4-tetrahydroisoquinolin-6-ol

[0279] The title compound was prepared by analogy to Example 48 except that 1-cyclohexyl-6-methoxy-2-phenyl-1,2,3,4-tetrahydroisoquinoline was used instead of 6-methoxy-1-(4-methoxy-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinoline, and hexanes:EtOAc 6:1 was used as eluent for flash chromatography purification.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  0.90-1.15 (6H, overlapping m), 1.60-1.70 (4H, overlapping m), 1.93 (1H, m), 2.90 (2H, m), 3.41 (1H, m), 3.66 (1H, m) 4.11 (1H, d, J 7.5 Hz), 4.98 (1H, broad s), 6.55-6.59 (2H, overlapping m), 6.65 (1H, t, J 7.0 Hz), 6.82 (2H, d, J 8.5 Hz), 6.90 (1H, d, J 8.0 Hz) and 7.17-7.23 (2H, overlapping m). MS 308 (M+1).

#### Example 51

##### 1-Isopropyl-6-methoxy-2-phenyl-1,2,3,4-tetrahydroisoquinoline

[0280] The title compound was prepared by analogy to Preparation 65 except that N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-isobutyramide was used instead of 4-methoxy-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide, and the final residue was not subjected to flash chromatography.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  0.92 (3H, d, J 6.5 Hz), 1.05 (3H, d, J 7.0 Hz), 2.11 (1H, m), 2.95 (2H, m), 3.42 (1H, m), 3.70 (1H, m), 3.76 (3H, s), 4.32 (1H, d, J 8.0 Hz) and 6.73-7.23 (8H, overlapping m). MS 282 (M+1).

#### Example 52

##### 1-Isopropyl-2-phenyl-1,2,3,4-tetrahydroisoquinolin-6-ol

[0281] The title compound was prepared by analogy to Example 48 except that 1-isopropyl-6-methoxy-2-phenyl-1,2,3,4-tetrahydroisoquinoline was used instead of 6-methoxy-1-(4-methoxy-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinoline, and hexanes:EtOAc 6:1 was used as eluent for flash chromatography purification.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  0.90 (3H, d, J 6.5 Hz), 1.02 (3H, d, J 7.0 Hz), 2.06 (1H, m), 2.90 (2H, m), 3.42 (1H, m), 3.66 (1H, m), 4.29 (1H, d, J 8.0 Hz), 4.77 (1H, s) and 6.56-7.23 (8H, overlapping m). MS 268 (M+1).

#### Example 53

##### 1,2-Diphenyl-1,2,3,4-tetrahydroisoquinolin-6-ol

[0282] The title compound was prepared by analogy to Example 48 except that 6-methoxy-1,2-diphenyl-1,2,3,4-tetrahydroisoquinoline was used instead of 6-methoxy-1-(4-methoxy-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinoline, and hexanes:EtOAc 6:1 was used as eluent for flash chromatography purification.

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  2.84 (2H, m), 3.45 (1H, m), 3.65 (1H, m), 4.98 (1H, broad s), 5.74 (1H, s), 6.56-7.24 (13H, overlapping m). MS 302 (M+1).

#### Example 54

##### 3-[4-(6-Methoxy-2-phenyl-1,2,3,4-tetrahydroisoquinolin-1-yl)-phenyl]-E-acrylic acid methyl/ethyl ester

[0283] The title compound was prepared by analogy to Preparation 65 except that 3-(4-[[2-(3-methoxy-phenyl)-ethyl]-phenyl-carbamoyl]-phenyl)-E-acrylic acid ethyl ester was used instead of 4-methoxy-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide. This compound was obtained as an approximately 1:1 mixture of methyl and ethyl esters (as determined by thin layer chromatography and mass spectrometry) due to transesterification.

MS 400 (R= Me, M+1) and 414 (R=Et, M+1).

**Example 55****-3-[4-(6-Hydroxy-2-phenyl-1,2,3,4-tetrahydro-isoquinolin-1-yl)-phenyl]-E-acrylic acid methyl/ethyl esters**

**[0284]** The title compound was prepared by analogy to Example 48 except that E-3 [4-(6-methoxy-2-phenyl-1,2,3,4-tetrahydro-isoquinolin-1-yl)-phenyl]-acrylic acid methyl/ethyl esters was used instead of 6-methoxy-1-(4-methoxy-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinoline, and hexanes:EtOAc 3:1 was used as the eluent for flash chromatography purification.

**Example 56****3-[4-(6-Hydroxy-2-phenyl-1,2,3,4-tetrahydroisoquinolin-1-yl)-phenyl]-E-acrylic acid methyl ester**

**[0285]** To 5 ml of the crude mixture of E-3-[4-(6-hydroxy-2-phenyl-1,2,3,4-tetrahydroisoquinolin-1-yl)-phenyl]-acrylic acid methyl/ethyl esters (0.045g) in MeOH (see Ex. 55) was added a catalytic amount of NaOMe (prepared by dissolving Na in MeOH) as a solution in MeOH (1 ml). The reaction was stirred at rt for 36 hours. The pH of the solution was adjusted to about 7 to 8 with 1N HCl and the product was extracted with CH<sub>2</sub>Cl<sub>2</sub>:EtOAc (4:1, 3x15 ml). The combined organics were dried over MgSO<sub>4</sub>, filtered, and concentrated *in vacuo*. The residue was subjected to flash chromatography (SiO<sub>2</sub>, hexanes:EtOAc 3:1) to give the desired product (0.012 g, 27% yield).

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 2.84 (2H, m), 3.44 (1H, m), 3.64 (1H, m), 3.76 (3H, s), 5.71 (1H, s), 6.34 (1H, d, J 16.0 Hz), 6.63-6.80 (5H, overlapping m) 7.08-7.35 (5H, overlapping m), 7.36 (2H, d, J 8.5 Hz) and 7.60 (1H, d, J 16.0 Hz). MS 386 (M+1).

**Example 57****3-[4-(6-Hydroxy-2-phenyl-1,2,3,4-tetrahydroisoquinolin-1-yl)-phenyl]-E-acrylic acid**

**[0286]** To a solution of 3-[4-(6-hydroxy-2-phenyl-1,2,3,4-tetrahydroisoquinolin-1-yl)-phenyl]-E-acrylic acid methyl/ethyl esters (0.045g, 0.115 mmol, 1.0 eq.) in THF (2 ml) was added NaOH (0.009g, 0.230 mmol, 2.0 eq.) as a solution in water (0.2 ml). Water (1.0 ml) was added and the reaction mixture was stirred at rt for 12 hours. The mixture was diluted with water (5 ml) and acidified to a pH of about 5. The aqueous mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>:EtOAc (4:1, 4x25 ml). The combined organics were dried over MgSO<sub>4</sub>, filtered, and concentrated *in vacuo*. The residue was subjected to flash chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>:MeOH 10:1) to give the desired compound (0.035 g, 0.0945 mmol, 82% yield).

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 2.87 (2H, m), 3.47 (1H, m), 3.68 (1H, m), 5.75 (1H, s), 6.37 (1H, d, J 16.0 Hz), 6.65-6.83 (5H, overlapping m), 7.13 (1H, d, J 8.0 Hz), 7.21-7.28 (4H, overlapping m), 7.42 (2H, d, J 8.5 Hz) and 7.72 (1H, d, J 16.0 Hz). MS 372 (M+H).

**Example 58****1-Benzyl-6-methoxy-2-phenyl-1,2,3,4-tetrahydroisoquinoline**

**[0287]** The title compound was prepared by analogy to Preparation 65 except that N-[2-(3-methoxy-phenyl)-ethyl]-2,N-diphenyl-acetamide was used instead of 4-methoxy-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide, and the final residue was not subjected to flash chromatography. MS 330 (M+1).

**Example 59****1-Benzyl-2-phenyl-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0288]** The title compound was prepared by analogy to Example 48 except that 1-benzyl-6-methoxy-2-phenyl-1,2,3,4-tetrahydroisoquinoline was used instead of 6-methoxy-1-(4-methoxy-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinoline, and hexanes:EtOAc 6:1 was used as eluent for flash chromatography purification.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 2.68 (1H, m), 2.67 (1H, m), 2.94 (1H, dd, 13.5 and 7.5 Hz), 3.21 (1H, dd, J 13.5 and 5.5 Hz), 3.49 (1H, m), 3.58 (1H, m), 4.83 (1H, dd, J 7.5 and 5.5 Hz), 5.18 (1H, broad s), 6.47-6.58 (3H, overlapping m), 6.74 (1H, t, J 7.5 Hz), 6.84 (2H, d, J 8.0 Hz), 7.00 (2H, dd, 7.0 and 1.5 Hz) and 7.17-7.26 (5H, overlapping m). MS 314 (M+1).



**Example 60****6-Methoxy-2-phenyl-1-thiophen-2-yl-1,2,3,4-tetrahydroisoquinoline**

**[0289]** The title compound was prepared by analogy to Preparation 65 except that thiophene-2-carboxylic acid [2-(3-methoxy-phenyl)-ethyl]-phenyl-amide was used instead of 4-methoxy-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide, and the final residue was not subjected to flash chromatography.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 2.90 (1H, m), 2.98 (1H, m), 3.54 (1H, m), 3.60 (1H, m), 3.80 (3H, s), 5.99 (1H, s) and 6.67-7.27 (11H overlapping m). MS 321 (M).

**Example 61****2-Phenyl-1-thiophen-2-yl-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0290]** The title compound was prepared by analogy to Example 48 except that 6-methoxy-2-phenyl-1-thiophen-2-yl-1,2,3,4-tetrahydroisoquinoline was used instead of 6-methoxy-1-(4-methoxy-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinoline, and hexanes:EtOAc from 4:1 to 1:1 was used as eluent for flash chromatography purification.

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 2.84 (1H, m), 2.91 (1H, m), 3.50 (1H, m), 3.56 (1H, m), 5.96 (1H, s), 6.61-6.67 (3H, overlapping m), 6.76-6.84 (2H overlapping m), 6.91 (2H, d J 7.5 Hz), 7.09 (2H, m) and 7.23 (2H, m). MS 308 (M+H).

**Example 62****6-Methoxy-1-naphthalen-2-yl-2-phenyl-1,2,3,4-tetrahydroisoquinoline**

**[0291]** The title compound was prepared by analogy to Preparation 65 except that naphthalene-2-carboxylic acid [2-(3-methoxy-phenyl)-ethyl]-phenyl-amide was used instead of 4-methoxy-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide, and the final residue was not subjected to flash chromatography.

MS (M) 365.

**Example 63****1-Naphthalen-2-yl-2-phenyl-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0292]** The title compound was prepared by analogy to Example 48 except that 6-methoxy-1-naphthalen-2-yl-2-phenyl-1,2,3,4-tetrahydroisoquinoline was used instead of 6-methoxy-1-(4-methoxy-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinoline, and neat CH<sub>2</sub>Cl<sub>2</sub> to 10% MeOH/CH<sub>2</sub>Cl<sub>2</sub> was used as eluent for flash chromatography purification.

<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ<sub>H</sub> 2.86 (2H, m), 3.44 (1H, m), 3.65 (1H, m), 5.87 (1H, s), 6.60-6.68 (3H, overlapping m), 6.87 (2H, d J 8.0 Hz), 7.12 (3H, m), 7.37 (3H, m), 7.51 (1H, s) and 3H, m). MS 352 (M+1).

**Example 64****6-Methoxy-2-phenyl-1-(3,4,5-trifluoro-phenyl)-1,2,3,4-tetrahydroisoquinoline**

**[0293]** The title compound was prepared by analogy to Preparation 65 except that 3,4,5-trifluoro-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide was used instead of 4-methoxy-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide, and the final residue was not subjected to flash chromatography.

MS 369 (M+1).

**Example 65****2-Phenyl-1-(3,4,5-trifluoro-phenyl)-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0294]** The title compound was prepared by analogy to Example 48 except that 6-methoxy-2-phenyl-1-(3,4,5-trifluoro-phenyl)-1,2,3,4-tetrahydroisoquinoline was used instead of 6-methoxy-1-(4-methoxy-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinoline, and neat CH<sub>2</sub>Cl<sub>2</sub> to 10% MeOH/CH<sub>2</sub>Cl<sub>2</sub> was used as eluent for flash chromatography purification.

<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ<sub>H</sub> 2.83 (2H, m), 3.38 (1H, m), 3.59 (1H, m), 5.70 (1H, s) and 6.61-7.18 (10H, overlapping m). MS 356 (M+1).

**Example 66****1-(4-Chloro-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0295]** The title compound was prepared by analogy to Example 48 except that 1-(4-chloro-phenyl)-6-methoxy-2-phenyl-1,2,3,4-tetrahydroisoquinoline was used instead of 6-methoxy-1-(4-methoxy-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinoline, and neat  $\text{CH}_2\text{Cl}_2$  to 10% MeOH/ $\text{CH}_2\text{Cl}_2$  was used as eluent for flash chromatography purification.  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  2.83 (2H, m), 3.37 (1H, m), 3.58 (1H, m), 5.71 (1H, s) and 6.62-7.18 (12H, overlapping m). MS 336/338 (M+1).

**Example 67****6-Methoxy-2-phenyl-1-thiazol-2-yl-1,2,3,4-tetrahydroisoquinoline**

**[0296]** The title compound was prepared by analogy to Preparation 65 except that thiazole-2-carboxylic acid [2-(3-methoxy-phenyl)-ethyl]-phenyl-amide was used instead of 4-methoxy-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenylbenzamide, and the final residue was not subjected to flash chromatography. MS 322 (M+1).

**Example 68****2-Phenyl-1-thiazol-2-yl-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0297]** The title compound was prepared by analogy to Example 48 except that 6-methoxy-2-phenyl-1-thiazol-2-yl-1,2,3,4-tetrahydroisoquinoline was used instead of 6-methoxy-1-(4-methoxy-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinoline, and neat  $\text{CH}_2\text{Cl}_2$  to 10% MeOH/ $\text{CH}_2\text{Cl}_2$  was used as eluent for flash chromatography purification.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  2.93 (1H, m), 3.01 (1H, m), 3.40 (1H, m), 3.77 (1H, m), 5.94 (1H, s), 6.59-6.86 (5H, overlapping m), 7.12 (1H, d J 3.0 Hz), 7.16-7.27 (3H, overlapping m) and 7.64 (1H, d J 3.0 Hz). MS 309 (M+1).

**Example 69****1-Adamantan-1-yl-6-methoxy-2-phenyl-1,2,3,4-tetrahydroisoquinoline**

**[0298]** The title compound was prepared by analogy to Preparation 65 except that adamantane-1-carboxylic acid [2-(3-methoxy-phenyl)-ethyl]-phenyl-amide was used instead of 4-methoxy-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenylbenzamide, and the final residue was subjected to flash chromatography using hexanes:EtOAc from 20:1 to 10:1. MS (M) 373.

**Example 70****1-Adamantan-1-yl-2-phenyl-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0299]** The title compound was prepared by analogy to Example 48 except that 1-adamantan-1-yl-6-methoxy-2-phenyl-1,2,3,4-tetrahydroisoquinoline was used instead of 6-methoxy-1-(4-methoxy-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinoline, and neat  $\text{CH}_2\text{Cl}_2$  to 10% MeOH/ $\text{CH}_2\text{Cl}_2$  was used as eluent for flash chromatography purification.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  1.54-2.10 (15H, overlapping m), 2.86 (1H, m), 3.06 (1H, m), 3.38 (1H, m), 3.84 (1H, m), 4.45 (1H, s) and 6.50-7.19 (8H, overlapping m). MS 360 (M+1).

**Example 71****6-Methoxy-2-phenyl-1-pyridin-4-yl-1,2,3,4-tetrahydroisoquinoline**

**[0300]** The title compound was prepared by analogy to Preparation 65 except that N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-isonicotinamide was used instead of 4-methoxy-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenylbenzamide, and the final residue was subjected to flash chromatography using hexanes:EtOAc from 4:1 to 2:1.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  2.82 (1H, m), 2.90 (1H, m), 3.47 (1H, m), 3.66 (1H, m), 3.79 (3H, s), 5.70 (1H, s), 5.70 (1H, s), 6.71-6.80 (5H, overlapping m), 7.18-7.25 (5H, overlapping m) and 8.46 (2H, m). MS 317 (M+1).

**Example 72****2-Phenyl-1-pyridin-4-yl-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0301]** The title compound was prepared by analogy to Example 48 except that 6-methoxy-2-phenyl-1-pyridin-4-yl-1,2,3,4-tetrahydroisoquinoline was used instead of 6-methoxy-1-(4-methoxy-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinoline, and hexanes:EtOAc from 4:1 to 1:1 was used as eluent for flash chromatography purification. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ<sub>H</sub> 2.84 (2H, overlapping m), 3.38 (1H, m), 3.65 (1H, m), 5.77 (1H, s), 6.60-6.71 (4H, overlapping m), 6.79 (2H, d, J 8.0 Hz), 7.15 (4H, m), 7.28 (2H, m) and 8.34 (1H, broad s). MS 303 (M+1).

**Example 73****1-(4-Iodo-phenyl)-6-methoxy-2-phenyl-1,2,3,4-tetrahydroisoquinoline**

**[0302]** The title compound was prepared by analogy to Preparation 65 except that 4-iodo-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide was used instead of 4-methoxy-N-[2-(3-methoxy-phenyl)-ethyl]-N-phenyl-benzamide, and the final residue was subjected to flash chromatography using a gradient from neat CH<sub>2</sub>Cl<sub>2</sub> to CH<sub>2</sub>Cl<sub>2</sub>:MeOH 60:1. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 2.83 (2H, m), 3.42 (1H, m), 3.62 (1H, m), 3.75 (3H, s), 5.65 (1H, s), 6.67-6.80 (5H, overlapping m), 6.94 (2H, d, J 8.5 Hz), 7.08-7.11 (3H, overlapping m) and 7.50 (2H, d, J 8.5 Hz). MS 442 (M+1).

**Example 74****1-(4-Iodo-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinolin-6-ol**

**[0303]** The title compound was prepared by analogy to Example 48 except that 1-(4-iodo-phenyl)-6-methoxy-2-phenyl-1,2,3,4-tetrahydroisoquinoline was used instead of 6-methoxy-1-(4-methoxy-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinoline, and CH<sub>2</sub>Cl<sub>2</sub>:MeOH in a gradient from 20:1 to 9:1 was used as eluent for flash chromatography purification. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 2.80 (2H, m), 3.43 (1H, m), 3.61 (1H, m), 5.65 (1H, s), 6.62-6.65 (2H, overlapping m), 6.74-6.81 (3H, overlapping m), 6.94 (2H, d, J 8.5 Hz), 7.05 (1H, m), 7.18-7.22 (2H, overlapping m) and 7.51 (2H, d, J 8.5 Hz). MS 428 (M+1).

**Example 75****E-3-[4-(6-Hydroxy-2-phenyl-1,2,3,4-tetrahydroisoquinolin-1-yl)-phenyl]-1-piperidin-1-yl-propenone**

**[0304]** To a stirred solution of piperidine (0.61 ml, 6.19 mmol, 1.0 eq.) and Et<sub>3</sub>N (2.58 ml, 18.6 mmol, 3.0 eq.) in toluene (10 ml) was added acryloyl chloride (1.14g, 12.4 mmol, 2.0 eq.) in a dropwise manner. The reaction was stirred overnight at RT. The reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> (20 ml) and washed sequentially with 1N HCl (1x 20 ml) and sat. NaHCO<sub>3</sub> (1x20 ml). The organic layer was dried over MgSO<sub>4</sub>, filtered, and concentrated in *vacuo*. The resulting material (1-piperidin-1-yl-propenone) was used without purification.

**[0305]** To a stirred solution of 1-(4-iodo-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinolin-6-ol (0.100g, 0.234 mmol, 1.0 eq.), 1-piperidin-1-yl-propenone (0.043 g, 0.351 mmol, 1.5 eq.), and Et<sub>3</sub>N (0.163 ml, 1.17 mmol, 5.0 eq.) in dry DMF (2 ml) under an atmosphere of N<sub>2</sub> was added Pd(PPh<sub>3</sub>)<sub>4</sub> (0.013g, 11.3 μmol, 0.05 eq.). The reaction mixture was heated to 100°C overnight. The reaction mixture was allowed to cool to RT, diluted with water (10 ml), and extracted with EtOAc (2x10ml). The combined organics were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated in *vacuo*. The residue was subjected to flash chromatography (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub> to CH<sub>2</sub>Cl<sub>2</sub>:MeOH 19:1) to give the desired compound. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 1.53-1.64 (6H, overlapping m), 2.77 (2H, m), 3.44-3.62 (6H, overlapping m), 5.68 (1H, s), 6.19-7.64 (14H, overlapping m). MS 439 (M+1).

**Example 76****E-3-[4-(6-Hydroxy-2-phenyl-1,2,3,4-tetrahydroisoquinolin-1-yl)-phenyl]-acrylamide**

**[0306]** To a stirred solution of 1-(4-iodo-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinolin-6-ol (0.100 g, 0.234 mmol), triethylamine (0.163 ml, 1.17 mmol), and acrylamide (0.082 g, 0.351 mmol) in DMF (2 ml), under an atmosphere of N<sub>2</sub>, was added Pd(PPh<sub>3</sub>)<sub>4</sub> (0.014 g, 0.012 mmol). The solution was heated to 100°C overnight, then cooled to RT and diluted with water (20 ml). The aqueous mixture was extracted with EtOAc (3x10 ml). The combined organics were dried over MgSO<sub>4</sub>, filtered, and concentrated in *vacuo*. The residue was subjected to flash chromatography (SiO<sub>2</sub>, 5%

MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to give the desired product (0.030 g, 0.089 mmol, 35% yield).

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 2.80 (2H, m), 3.19 (1H, m), 3.42 (1H, m), 5.67 (1H, s), 5.88 (1H, br s), 6.17 (1H, br s), 6.38 (1H, d, J 15.5 Hz), 6.67-6.78 (5H, overlapping m), 7.04-7.32 (7H, overlapping m) and 7.52 (1H, d, 15.5 Hz). MS 371 (M+1).

#### Example 77

##### E-3-[4-(6-Hydroxy-2-phenyl-1,2,3,4-tetrahydroisoquinolin-1-yl)-phenyl]-1

##### morpholin-4-yl-propenone

[0307] The title compound was prepared by analogy to Example 75 except that morpholine was used instead piperidine,

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 3.10 (2H, m), 3.56-3.74 (10H, overlapping m), 5.71 (1H, s), 6.49-7.67 (14H, overlapping m). MS 441 (M+1).

#### Example 78

##### E-3-[4-(6-Hydroxy-2-phenyl-1,2,3,4-tetrahydroisoquinolin-1-yl)-phenyl]-N,N

##### dimethyl-acrylamide

[0308] The title compound was prepared by analogy to Example 75 except that dimethylamine was used instead piperidine.

MS 399 (M+1).

#### Example 79

##### 2,2,2-Trifluoro-1-[6-methoxy-1-[1-(1-methyl-1H-imidazole-4-sulfonyl)-piperidin-4-yl]-3,4-dihydro-1H-isoquinolin-2-yl]-ethanone

[0309] To a stirred solution of 6-methoxy-1-[1-(1-methyl-1H-imidazole-4-sulfonyl)-piperidin-4-yl]-1,2,3,4-tetrahydroisoquinoline (0.049 g, 0.126 mmol) and triethylamine (0.135 g, 0.132 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (5 ml) cooled to 0°C under an atmosphere of N<sub>2</sub> was added trifluoroacetic acid anhydride (0.028 g, 0.132 mmol). The reaction mixture was stirred overnight, slowly warming to RT. The reaction was quenched by the addition of water (1ml) and diluted with CH<sub>2</sub>Cl<sub>2</sub> (5ml). The layers were separated and the organic layer was further diluted with CH<sub>2</sub>Cl<sub>2</sub> (15 ml), washed with sat. NaHCO<sub>3</sub> (2x 3ml) and brine (1x3 ml), dried over MgSO<sub>4</sub>, filtered, and concentrated in *vacuo*. The residue was purified by radial chromatography (SiO<sub>2</sub>, 1mm CH<sub>2</sub>Cl<sub>2</sub> to 2% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) to give the desired product as an oil (0.043 g, 0.088 mmol, 71% yield).

<sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub> 1.41-1.80 (5H, overlapping m), 2.53 (2H, m), 2.92 (2H, m), 3.68 (1H, m), 3.73 (3H, s), 3.77 (3H, s), 3.90 (3H, m), 5.23 (1H, d, J 8.0 Hz), 6.65-7.00 (3H, m), 7.39 (1H, d, J 1.5Hz), and 7.45 (1H, d, J 1.5 Hz).

#### Example 80

##### 2,2,2-Trifluoro-1-[6-hydroxy-1-[1-(1-methyl-1H-imidazole-4-sulfonyl)-piperidin-4-yl]-3,4-dihydro-1H-isoquinolin-2-yl]-ethanone

[0310] To a stirred solution of 2,2,2-trifluoro-1-[6-methoxy-1-[1-(1-methyl-1H-imidazole-4-sulfonyl)-piperidin-4-yl]-3,4-dihydro-1H-isoquinolin-2-yl]-ethanone (0.036 g, 0.074 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3 ml) under an atmosphere of N<sub>2</sub> cooled to -78°C was added boron tribromide as a 1.0 M solution in CH<sub>2</sub>Cl<sub>2</sub> (0.3 ml, 0.3 mmol). The reaction mixture was stirred overnight slowly, while allowing it to warm to RT. MeOH (1 ml) was added to the reaction mixture and stirring was continued at RT for 10 minutes. Sat. NaHCO<sub>3</sub> was added to the reaction mixture until the pH of the aqueous layer was 7. The mixture was then extracted with EtOAc (3x20 ml), the combined organics were washed with brine (1x20 ml), dried over MgSO<sub>4</sub>, filtered, and concentrated in *vacuo* to give the desired product as an off white solid (0.028 g, 0.059 mmol, 80% yield).

MS 473 (M+1).

**Example 81****2-Benzyl-7-methoxy-4-(4-methoxy-phenyl)-1,2,3,4-tetrahydroisoquinoline**

[0311] To a solution of 2-[benzyl-(3-methoxy-benzyl)-amino]-1-(4-methoxy-phenyl)-ethanol (16.23 g, 43.0 mmol, see Preparation 63) in  $\text{CH}_2\text{Cl}_2$  (280 ml) was added TFA (40.0 g, 27.0 ml, 350 mmol). The mixture was refluxed overnight. The reaction mixture was allowed to cool to RT and poured onto ice-water containing sat.  $\text{NaHCO}_3$  (250 ml). The mixture was extracted with  $\text{CH}_2\text{Cl}_2$  (3x150 ml) and the combined organics were dried over  $\text{MgSO}_4$ , filtered, and concentrated *in vacuo*. Flash chromatography ( $\text{SiO}_2$ , gradient elution 10-40 % EtOAc/Hexanes) of the residue afforded the title product (7.67 g, 21.3 mmol, 50% yield).

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  2.60 (1H, m), 3.03 (1H, m), 3.67 (2H, m), 3.77 (3H, s), 4.13 (1H, m), 6.54 (1H, d, J 2.5 Hz), 6.62 (1H, dd, J 8.5 and 2.5 Hz), 6.78 (3H, m), 7.08 (2H, d, J 8.5 Hz) and 7.73-7.28 (5H, m). MS 360 (M+1).

**Example 82****2,2,2-Trifluoro-1-[7-methoxy-4-(4-methoxy-phenyl)-3,4-dihydro-1H-isoquinolin-2-yl]-ethanone**

[0312] To a solution of 7-methoxy-4-(4-methoxy-phenyl)-1,2,3,4-tetrahydroisoquinoline (1.43 g, 5.31 mmol, see Preparation 64) and  $\text{Et}_3\text{N}$  (1.08 g, 10.6 mmol) in  $\text{CH}_2\text{Cl}_2$  (50 ml) cooled to  $0^\circ\text{C}$ , was added trifluoroacetic anhydride (1.89 g, 1.27 ml, 9.03 mmol). The reaction was stirred at  $0^\circ\text{C}$  for 4 hours, then the reaction mixture was poured into sat.  $\text{NaHCO}_3$  (50 ml). The mixture was extracted with  $\text{CH}_2\text{Cl}_2$  (3x50 ml). The organics were combined, dried over  $\text{MgSO}_4$ , filtered, and concentrated *in vacuo*. Flash chromatography (Biotage,  $\text{SiO}_2$ , 10% EtOAc/hexanes) gave the title product (1.74 g, 4.76 mmol, 90% yield).

$^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta_{\text{H}}$  3.69-3.75 (1H, m), 3.78 (3H, s), 3.79 (3H, s), 4.00-4.18 (2H, m), 4.72-4.99 (2H, m), 6.65-6.75 (2H, m), 6.81-6.87 (3H, m) and 6.98-7.02 (2H, m). MS 366 (M+1).

**Example 83****2,2,2-Trifluoro-1-[7-hydroxy-4-(4-hydroxy-phenyl)-3,4-dihydro-1H-isoquinolin-2-yl]-ethanone**

[0313] To a solution of 2,2,2-trifluoro-1-[7-methoxy-4-(4-methoxy-phenyl)-3,4-dihydro-1H-isoquinolin-2-yl]-ethanone (1.74 g, 4.76 mmol, see Example 82) in  $\text{CH}_2\text{Cl}_2$  (100 ml) cooled to  $0^\circ\text{C}$  was added  $\text{BBr}_3$  as a 1.0 M solution in  $\text{CH}_2\text{Cl}_2$  (14.28 ml, 14.28 mmol) in a dropwise manner. The reaction mixture was stirred at  $0^\circ\text{C}$  for 60 minutes, the ice bath removed, and stirring continued at RT for 90 minutes. MeOH (10 ml) was added to the reaction and stirring was continued at RT for 10 minutes. The reaction mixture was concentrated *in vacuo* and purified by flash chromatography (Biotage,  $\text{SiO}_2$ , 25-40% EtOAc/hexanes) to give the title product (1.36 g, 4.03 mmol, 85% yield).

$^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  3.77-4.13 (3H, m), 4.72-4.86 (2H, m), 6.59-6.76 (5H, m) and 6.86-6.90 (2H, m). MS 338 (M+1).

**Example 84****Cyclohexyl-[7-hydroxy-4-(4-hydroxy-phenyl)-3,4-dihydro-1H-isoquinolin-2-yl]-methanone**

[0314] A solution of 7-methoxy-4-(4-methoxy-phenyl)-1,2,3,4-tetrahydroisoquinoline (0.103 g, 0.382 mmol, 1.0 eq., see Preparation 64) and triethylamine (0.107 ml, 0.764 mmol, 2.0 eq.) in  $\text{CH}_2\text{Cl}_2$  (2 ml) was added to a vial containing cyclohexanecarbonyl chloride (0.084 g, 0.573 mmol, 1.5 eq.). The reaction mixture was stirred overnight at RT. To this mixture was added  $\text{BBr}_3$  as a 1.0 M solution in  $\text{CH}_2\text{Cl}_2$  (1.15 ml, 1.15 mmol, 3.0 eq.). The reaction mixture was stirred overnight at RT, then cooled to  $-78^\circ\text{C}$  and quenched with MeOH (1.5 ml). The reaction mixture was evaporated under a stream of nitrogen and purified by reverse phase HPLC (gradient elution 98:2  $\text{H}_2\text{O}$ :0.1% TFA, then 2:98 MeCN: $\text{H}_2\text{O}$ ). The exact yield was not determined.

$^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta_{\text{H}}$  1.12-1.94 (10H, overlapping m), 2.67 (1H, m), 3.77 (1H, m), 3.94 (1H, m), 4.09 (1H, m), 4.35 (1H, d, J 17.5 Hz), 5.06 (1H, d, J 17.5 Hz) and 6.58-6.88 (7H, overlapping m). MS 352 (M+1).

**Example 85****[7-Hydroxy-4-(4-hydroxy-phenyl)-3,4-dihydro-1H-isoquinolin-2-yl]-phenyl-methanone**

[0315] The title compound was prepared by analogy to Example 84 except benzoyl chloride was used in place of

cyclohexanecarbonyl chloride.

<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ<sub>H</sub> 3.68 (2H, m), 3.84-4.18 (3H, overlapping m), 4.52 (1H, d, J 17.5 Hz), 5.22 (1H, d, J 17.5 Hz), 6.57-6.96 (8H, overlapping m) and 7.19-7.46 (4H, m). MS 346 (M+1).

## Example 86

### 2-Benzenesulfonyl-4-(4-hydroxy-phenyl)-1,2,3,4-tetrahydroisoquinolin-7-ol

[0316] A solution of 7-methoxy-4-(4-methoxy-phenyl)-1,2,3,4-tetrahydroisoquinoline (0.103 g, 0.382 mmol, 1.0 eq., see Preparation 64) and triethylamine (0.107 ml, 0.764 mmol, 2.0 eq.) in CH<sub>2</sub>Cl<sub>2</sub> (2 ml) was added to a vial containing benzenesulfonyl chloride (0.101 g, 0.573 mmol, 1.5 eq.). The reaction mixture was stirred overnight at RT. To this mixture was added BBr<sub>3</sub> as a 1.0M solution in CH<sub>2</sub>Cl<sub>2</sub> (1.15 ml, 1.15 mmol, 3.0 eq.). The reaction mixture was again stirred overnight at RT, then cooled to -78°C and quenched with MeOH (1.5 ml). The reaction mixture was evaporated under a stream of nitrogen and purified by reverse phase HPLC (gradient elution 98:2 H<sub>2</sub>O:0.1% TFA, then 2:98 MeCN: H<sub>2</sub>O). The exact yield was not determined.

<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ<sub>H</sub> 3.04 (1H, dd, J 12.0 and 7.5 Hz), 3.55 (1H, dd, J 12.0 and 5.0 Hz), 4.02 (1H, dd, J 6.5 and 6.0 Hz), 4.12 (1H, d, J 15.0 Hz), 4.27 (1H, d, J 15.0 Hz), 6.52 (2H, m), 6.65 (3H, m), 6.87 (1H, d, J 8.5 Hz), 7.55 (2H, m), 7.56 (1H, m) and 7.75 (1H, d, J 8.0 Hz). MS 382 (M+1).

## Example 87

### 4-(4-Hydroxy-phenyl)-2-(naphthalene-1-sulfonyl)-1,2,3,4-tetrahydroisoquinolin-7-ol

[0317] The title compound was prepared by analogy to Example 86 except naphthalene-1-sulfonyl chloride was used in place of benzenesulfonyl chloride.

<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ<sub>H</sub> 3.34 (1H, dd, J 12.0 and 7.0 Hz), 3.64 (1H, dd, J 12.0 and 4.5 Hz), 3.93 (1H, t, J 6.0 Hz), 4.36 (1H, d, J 15.5 Hz), 4.42 (1H, d, J 15.5 Hz), 6.45-6.68 (7H, overlapping m), 7.93 (1H, m), 8.10-8.18 (2H, overlapping m) and 8.54 (1H, m). MS 432 (M+1).

## Example 88

### 2-(Biphenyl-4-sulfonyl)-4-(4-hydroxy-phenyl)-1,2,3,4-tetrahydroisoquinolin-7-ol

[0318] The title compound was prepared by analogy to Example 86 except biphenyl-4-sulfonyl chloride was used in place of benzenesulfonyl chloride.

<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ<sub>H</sub> 3.09 (1H, dd, J 12.0 and 7.5 Hz), 3.59 (1H, dd, J 12.0 and 5.0 Hz), 4.05 (1H, dd, J 7.5 and 5.0 Hz), 4.17 (1H, d, J 15.0 Hz), 4.32 (1H, J 15.0 Hz), 6.54 (3H, m), 6.64 (3H, m), 6.88 (2H, m), 7.36-7.47 (3H, overlapping m), 7.66 (2H, m) and 7.80 (2H, m). MS 458 (M+1).

## Example 89


### 4-(4-Hydroxy-phenyl)-2-phenylmethanesulfonyl-1,2,3,4-tetrahydroisoquinolin-7-ol

[0319] The title compound was prepared by analogy to Example 86 except phenyl-methanesulfonyl chloride was used in place of benzenesulfonyl chloride.

<sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ<sub>H</sub> 3.19 (1H, m), 3.56 (1H, dd, 12.5 and 5.0 Hz), 3.88 (1H, J 7.5 and 6.0 Hz), 4.20-4.39 (4H, overlapping m), 6.48-6.58 (2H, overlapping m), 6.52-6.71 (3H, overlapping m), 6.85 (2H, m) and 7.22-7.38 (5H, overlapping m). MS 396 (M+1).

## Claims

1. A compound of the formula:



wherein:

A<sup>1</sup> is hydrogen, hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, or (C<sub>1</sub>-C<sub>4</sub>)alkanoyloxy, said (C<sub>1</sub>-C<sub>4</sub>)alkoxy or said (C<sub>1</sub>-C<sub>4</sub>)alkanoyloxy being optionally substituted by hydroxy, halo, or a partially saturated, fully saturated, or fully unsaturated five to twelve membered ring optionally having up to four heteroatoms independently selected from oxygen, sulfur, and nitrogen, or A<sup>1</sup> is R<sup>3</sup>-(C<sub>1</sub>-C<sub>4</sub>)alkoxy wherein R<sup>3</sup> is pyrrolidino, piperidino, morpholino, or dimethylamino; A<sup>2</sup>, A<sup>3</sup>, and A<sup>4</sup> are independently selected from hydrogen, hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, and halo; R<sup>1</sup> is phenyl; pyridyl; piperidinyl; (C<sub>1</sub>-C<sub>7</sub>)alkyl; adamantyl; a partially saturated, fully saturated, or fully unsaturated three to twelve membered ring optionally having up to four heteroatoms selected independently from oxygen, sulfur, and nitrogen; a bicyclic ring consisting of two fused independently partially saturated, fully saturated, or fully unsaturated five to six membered rings, wherein said bicyclic ring includes up to four heteroatoms independently selected from oxygen, sulfur, and nitrogen; or a bicyclic ring system consisting of two rings joined by a covalent bond, said rings being independently partially saturated, fully saturated, or fully unsaturated three to eight membered rings, wherein said bicyclic ring system includes up to four heteroatoms independently selected from oxygen, sulfur, and nitrogen; wherein each of the above R<sup>1</sup> groups is optionally substituted with up to seven fluoro atoms, or with up to three substituents independently selected from Group A, wherein Group A consists of hydroxy, halo, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, R<sup>3</sup>-(C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>2</sub>-C<sub>4</sub>)alkenyl-COOR<sup>7</sup> wherein R<sup>7</sup> is hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-COOR<sup>7</sup>, (C<sub>1</sub>-C<sub>4</sub>)alkanoyloxy-(C<sub>2</sub>-C<sub>4</sub>)alkenyl, (C<sub>2</sub>-C<sub>4</sub>)alkenyl-CONR<sup>4</sup>R<sup>5</sup> wherein R<sup>4</sup> and R<sup>5</sup> are independently hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkylene, or (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, or R<sup>4</sup> and R<sup>5</sup> taken together with the nitrogen atom to which they are attached form pyrrolidino, piperidino, morpholino, or hexamethyleneimino, (C<sub>0</sub>-C<sub>4</sub>)alkyl-CONR<sup>4</sup>R<sup>5</sup>, (C<sub>0</sub>-C<sub>4</sub>)alkyl-NR<sup>4</sup>R<sup>5</sup>, OCH<sub>2</sub>CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup> wherein R<sup>8</sup> and R<sup>9</sup> are independently methyl or ethyl, or R<sup>8</sup> and R<sup>9</sup> taken together with the nitrogen atom to which they are attached form pyrrolidino, piperidino, morpholino, or hexamethyleneimino, propyl-R<sup>8</sup>R<sup>9</sup>, and SO<sub>2</sub>-R<sup>6</sup> wherein R<sup>6</sup> is imidazolyl, thienyl, benzathienyl, or isoxazolyl, optionally substituted with up to three substituents independently selected from (C<sub>1</sub>-C<sub>4</sub>)alkyl;

X is a covalent bond, (CH<sub>2</sub>)<sub>n</sub> where n is 1, 2, or 3, (C<sub>0</sub>-C<sub>1</sub>)alkylene-phenylene-(C<sub>0</sub>-C<sub>1</sub>)alkylene, CO<sub>2</sub>, (C<sub>0</sub>-C<sub>3</sub>)alkylene-CO-(C<sub>0</sub>-C<sub>3</sub>)alkylene, or (C<sub>0</sub>-C<sub>4</sub>)alkylene-SO<sub>2</sub>-(C<sub>0</sub>-C<sub>4</sub>)alkylene;

R<sup>2</sup> is (C<sub>1</sub>-C<sub>9</sub>)alkyl; (C<sub>2</sub>-C<sub>4</sub>)alkenyl; benzhydryl; a partially saturated, fully saturated, or fully unsaturated three to eight membered ring optionally having up to four heteroatoms selected independently from oxygen, sulfur, and nitrogen; a bicyclic ring consisting of two fused independently partially saturated, fully saturated, or fully unsaturated five to six membered rings, wherein said bicyclic ring includes up to four heteroatoms independently selected from oxygen, sulfur, and nitrogen; or a bicyclic ring system consisting of two rings joined by a covalent bond, said rings being independently partially saturated, fully saturated, or fully unsaturated three to eight membered rings, wherein said bicyclic ring system includes up to four heteroatoms independently selected from oxygen, sulfur, and nitrogen; wherein said (C<sub>1</sub>-C<sub>9</sub>)alkyl is optionally substituted with one to seven fluoro substituents, or up to three substituents independently selected from Group B, wherein Group B consists of chloro, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, amino, and (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonyl; wherein said (C<sub>2</sub>-C<sub>4</sub>)alkenyl is optionally substituted with up to three substituents independently selected from Group C, wherein Group C consists of halo, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, amino, and (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonyl; and wherein said benzhydryl, said 5 to 8 membered ring, said bicyclic ring, and said bicyclic ring system is optionally substituted with up to three substituents independently selected from Group D, wherein Group D consists of halo, hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, imidazolyl, amino, (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonylamino, and (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonyl; and

p is 0, 1, or 2;

with the proviso that

when X is (CH<sub>2</sub>)<sub>2</sub> or (CH<sub>2</sub>)<sub>3</sub>, p is 0, and R<sup>1</sup> is phenyl or phenyl substituted with a single chloro, fluoro, bromo, hydroxy, methoxy, pyrrolidinoethoxy, piperidinoethoxy, or morpholinoethoxy substituent, then R<sup>2</sup> is not phenyl, methoxyphenyl, tert-butyl, or cyclopentyl;  
 when X is CH<sub>2</sub>, (CH<sub>2</sub>)<sub>2</sub>, COCH<sub>2</sub>, or CH<sub>2</sub>CO, A<sup>1</sup> is hydrogen, and R<sup>1</sup> is phenyl, then R<sup>2</sup> is not phenyl; and  
 when X is a covalent bond, p is 0, A<sup>1</sup> is hydrogen or methoxy, and R<sup>1</sup> is phenyl or phenyl substituted with a single chloro, fluoro, bromo, methoxy, pyrrolidinoethoxy, or piperidinoethoxy substituent, then R<sup>2</sup> is not phenyl or *m*-fluorophenyl.

2. A compound of claim 1 wherein:

A<sup>1</sup> is hydroxy;  
 A<sup>2</sup>, A<sup>3</sup>, and A<sup>4</sup> are hydrogen; and  
 p is 0.

3. A compound of claim 1 wherein:

A<sup>1</sup> is hydroxy;  
 A<sup>2</sup>, A<sup>3</sup>, and A<sup>4</sup> are hydrogen;  
 p is 0;  
 R<sup>1</sup> is phenyl, cyclohexyl, pyridyl, thienyl, isopropyl, or adamantyl; wherein each of said R<sup>1</sup> groups is optionally substituted with up to three fluoro atoms, or with one substituent selected from iodo, chloro, bromo, hydroxy, methoxy, pyrrolidino-ethoxy, dimethylamino, COOR<sup>7</sup> wherein R<sup>7</sup> is hydrogen or methyl, or ethenyl-CONR<sup>4</sup>R<sup>5</sup> wherein R<sup>4</sup> and R<sup>5</sup> are both methyl, or R<sup>4</sup> and R<sup>5</sup> taken together with the nitrogen atom to which they are attached form piperidino or morpholino;  
 X is a covalent bond, CH<sub>2</sub>, CH<sub>2</sub>-phenylene, CO<sub>2</sub>, CO-(C<sub>0</sub>-C<sub>2</sub>)alkylene, or SO<sub>2</sub>-(C<sub>0</sub>-C<sub>2</sub>)alkylene; and  
 R<sup>2</sup> is methyl, t-butyl, phenyl, cyclohexyl, isoxazolyl, tetrahydropyranyl, naphthyl, or benzodioxolyl; wherein each of said methyl or t-butyl is optionally substituted with one to three fluoro substituents, or up to two substituents independently selected from amino and methylcarbonyl; and wherein each of said phenyl, cyclohexyl, isoxazolyl, tetrahydropyranyl, naphthyl, or benzodioxolyl is optionally substituted with up to three fluoro substituents, or up to two substituents independently selected from hydroxy, (C<sub>1</sub>-C<sub>3</sub>)alkyl, amino, and methylcarbonyl.

4. A compound of claim 1 wherein:

said compound is of formula (I);  
 A<sup>1</sup> is hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>1</sub>-C<sub>4</sub>)alkanoyloxy, or pyrrolidino-ethoxy;  
 A<sup>2</sup>, A<sup>3</sup>, and A<sup>4</sup> are hydrogen;  
 p is 0 or 1;  
 R<sup>1</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>4</sub>-C<sub>7</sub>)cycloalkyl, adamantyl, phenyl, pyridyl, or thienyl, wherein each of said phenyl, pyridyl, or thienyl groups is optionally substituted with up to three fluoro atoms, or with one substituent selected from iodo, chloro, bromo, hydroxy, methoxy, dimethylamino, OCH<sub>2</sub>CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, COOR<sup>7</sup>, ethenyl-COOR<sup>7</sup>, or ethenyl-CONR<sup>4</sup>R<sup>5</sup> wherein R<sup>4</sup> and R<sup>5</sup> are both methyl, or R<sup>4</sup> and R<sup>5</sup> taken together with the nitrogen atom to which they are attached form pyrrolidino, piperidino, hexamethyleneimino, or morpholino;  
 X is a covalent bond, CH<sub>2</sub>, CH<sub>2</sub>-phenylene, CO<sub>2</sub>, CO-(C<sub>0</sub>-C<sub>2</sub>)alkylene, or SO<sub>2</sub>-(C<sub>0</sub>-C<sub>2</sub>)alkylene; and  
 R<sup>2</sup> is (C<sub>1</sub>-C<sub>7</sub>)alkyl, phenyl, benzyl, thienyl, (C<sub>5</sub>-C<sub>7</sub>)cycloalkyl, isoxazolyl, imidazolyl, tetrahydropyranyl, naphthyl, or benzodioxolyl, wherein said (C<sub>1</sub>-C<sub>7</sub>)alkyl is optionally substituted with one to three fluoro substituents, or up to two substituents independently selected from amino and methylcarbonyl, and wherein each of said phenyl, thienyl, (C<sub>5</sub>-C<sub>7</sub>)cycloalkyl, isoxazolyl, tetrahydropyranyl, naphthyl, and benzodioxolyl is optionally substituted with up to three fluoro substituents, or up to two substituents independently selected from hydroxy, methoxy, (C<sub>1</sub>-C<sub>3</sub>)alkyl, amino, and methylcarbonyl.

5. A compound of claim 1 wherein:

said compound is of formula (II);  
 A<sup>1</sup> is hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, or (C<sub>1</sub>-C<sub>4</sub>)alkanoyloxy;  
 A<sup>2</sup>, A<sup>3</sup>, and A<sup>4</sup> are hydrogen;



R<sup>1</sup> is (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>4</sub>-C<sub>7</sub>)cycloalkyl, adamantyl, phenyl, pyridyl, or thienyl, wherein each of said phenyl, pyridyl, thienyl, or (C<sub>5</sub>-C<sub>7</sub>)cycloalkyl groups is optionally substituted with up to three fluoro atoms, or with one substituent selected from iodo, chloro, bromo, hydroxy, methoxy, dimethylamino, OCH<sub>2</sub>CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, COOR<sup>7</sup>, or ethenyl-CONR<sup>4</sup>R<sup>5</sup> wherein R<sup>4</sup> and R<sup>5</sup> are both methyl, or R<sup>4</sup> and R<sup>5</sup> taken together with the nitrogen atom to which they are attached form pyrrolidino, piperidino, hexamethyleneimino, or morpholino;


X is a covalent bond, CH<sub>2</sub>, CH<sub>2</sub>-phenylene, CO<sub>2</sub>, CO-(C<sub>0</sub>-C<sub>2</sub>)alkylene, or SO<sub>2</sub>-(C<sub>0</sub>-C<sub>2</sub>)alkylene; and

R<sup>2</sup> is (C<sub>1</sub>-C<sub>7</sub>)alkyl, phenyl, benzyl, thienyl, (C<sub>5</sub>-C<sub>7</sub>)cycloalkyl, isoxazolyl, tetrahydropyranyl, naphthyl, or benzodioxolyl, wherein said (C<sub>1</sub>-C<sub>7</sub>)alkyl is optionally substituted with one to three fluoro atoms, or up to two substituents independently selected from amino and methylcarbonyl, and wherein each of said phenyl, thienyl, cyclohexyl, isoxazolyl, tetrahydropyranyl, naphthyl, and benzodioxolyl is optionally substituted with up to three fluoro atoms, or up to two substituents independently selected from hydroxy, methoxy, and (C<sub>1</sub>-C<sub>3</sub>)alkyl.

6. A compound of claim 1 which is:

2,2,2-trifluoro-1-[7-hydroxy-4-(4-hydroxy-phenyl)-3,4-dihydro-1H-isoquinolin-2-yl]-ethanone;  
2,2,2-trifluoro-1-[6-hydroxy-1-(4-hydroxyphenyl)-3,4-dihydro-1H-isoquinolin-2-yl]-ethanone;  
2,2,2-trifluoro-1-[6-hydroxy-1(R)-(4-hydroxyphenyl)-3,4-dihydro-1H-isoquinolin-2-yl]-ethanone;  
2-benzyl-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol;  
2-benzenesulfonyl-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol;  
2-(4-isopropylbenzenesulfonyl)-1-[4-(2-pyrrolidin-1-yl-ethoxy)phenyl]-1,2,3,4-tetrahydroisoquinolin-6-ol;  
1-(4-hydroxy-phenyl)-2-phenyl-1,2,3,4-tetrahydroisoquinolin-6-ol;  
3-[4-(6-hydroxy-2-phenyl-1,2,3,4-tetrahydroisoquinolin-1-yl)-phenyl]-1-piperidin-1-yl-propenone;  
3-[4-(6-hydroxy-2-phenyl-1,2,3,4-tetrahydroisoquinolin-1-yl)-phenyl]-1-morpholin-4-yl-propenone; or  
3-[4-(6-hydroxy-2-phenyl-1,2,3,4-tetrahydroisoquinolin-1-yl)-phenyl]-N,N-dimethyl-acrylamide.

7. A compound of the formula:



A, wherein Group A consists of hydroxy, halo, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, R<sup>3</sup>-(C<sub>1</sub>-C<sub>4</sub>)alkoxy, (C<sub>2</sub>-C<sub>4</sub>)alkenyl-COOR<sup>7</sup> wherein R<sup>7</sup> is hydrogen or (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>0</sub>-C<sub>4</sub>)alkyl-COOR<sup>7</sup>, (C<sub>1</sub>-C<sub>4</sub>)alkanoyloxy-(C<sub>2</sub>-C<sub>4</sub>)alkenyl, (C<sub>2</sub>-C<sub>4</sub>)alkenyl-CONR<sup>4</sup>R<sup>5</sup> wherein R<sup>4</sup> and R<sup>5</sup> are independently hydrogen, (C<sub>1</sub>-C<sub>4</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy-(C<sub>1</sub>-C<sub>4</sub>)alkylene, or (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, or R<sup>4</sup> and R<sup>5</sup> taken together with the nitrogen atom to which they are attached form pyrrolidino, piperidino, morpholino, or hexamethyleneimino, (C<sub>0</sub>-C<sub>4</sub>)alkyl-CONR<sup>4</sup>R<sup>5</sup>, (C<sub>0</sub>-C<sub>4</sub>)alkyl-NR<sup>4</sup>R<sup>5</sup>, OCH<sub>2</sub>CH<sub>2</sub>NR<sup>8</sup>R<sup>9</sup> wherein R<sup>8</sup> and R<sup>9</sup> are independently methyl or ethyl, or R<sup>8</sup> and R<sup>9</sup> taken together with the nitrogen atom to which they are attached form pyrrolidino, piperidino, morpholino, or hexamethyleneimino, propyl-R<sup>8</sup>R<sup>9</sup>, and SO<sub>2</sub>-R<sup>6</sup> wherein R<sup>6</sup> is imidazolyl, thienyl, benzathienyl, or isoxazolyl, optionally substituted with up to three substituents independently selected from (C<sub>1</sub>-C<sub>4</sub>)alkyl;

X is (C<sub>0</sub>-C<sub>1</sub>)alkylene-phenylene-(C<sub>0</sub>-C<sub>1</sub>)alkylene, CO<sub>2</sub>, CO, (C<sub>1</sub>-C<sub>3</sub>)alkylene-CO-(C<sub>1</sub>-C<sub>3</sub>)alkylene, (C<sub>0</sub>-C<sub>3</sub>)alkylene-CO-(C<sub>2</sub>-C<sub>3</sub>)alkylene, (C<sub>2</sub>-C<sub>3</sub>)alkylene-CO-(C<sub>0</sub>-C<sub>3</sub>)alkylene, or (C<sub>0</sub>-C<sub>4</sub>)alkylene-SO<sub>2</sub>-(C<sub>0</sub>-C<sub>4</sub>)alkylene;

R<sup>2</sup> is (C<sub>1</sub>-C<sub>9</sub>)alkyl; (C<sub>2</sub>-C<sub>4</sub>)alkenyl; benzhydryl; a partially saturated, fully saturated, or fully unsaturated three to eight membered ring optionally having up to four heteroatoms selected independently from oxygen, sulfur, and nitrogen; a bicyclic ring consisting of two fused independently partially saturated, fully saturated, or fully unsaturated five to six membered rings, wherein said bicyclic ring includes up to four heteroatoms independently selected from oxygen, sulfur, and nitrogen; or a bicyclic ring system consisting of two rings joined by a covalent bond, said rings being independently partially saturated, fully saturated, or fully unsaturated three to eight membered rings, wherein said bicyclic ring system includes up to four heteroatoms independently selected from oxygen, sulfur, and nitrogen; wherein said (C<sub>1</sub>-C<sub>9</sub>)alkyl is optionally substituted with one to seven fluoro substituents, or up to three substituents independently selected from Group B, wherein Group B consists of chloro, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, amino, and (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonyl; wherein said (C<sub>2</sub>-C<sub>4</sub>)alkenyl is optionally substituted with up to three substituents independently selected from Group C, wherein Group C consists of halo, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, amino, and (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonyl; and wherein said benzhydryl, said 5 to 8 membered ring, said bicyclic ring, and said bicyclic ring system is optionally substituted with up to three substituents independently selected from Group D, wherein Group D consists of halo, hydroxy, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, imidazolyl, amino, (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonylamino, and (C<sub>1</sub>-C<sub>4</sub>)alkylcarbonyl; and p is 0, 1, or 2.

8. A method for treating or preventing a disease, disorder, condition, or symptom mediated by an estrogen receptor and/or caused by lowered estrogen level in a mammal, said method comprising administering to said mammal a therapeutically effective amount of a compound of claim 1.
9. A method for treating or preventing a disease, disorder, condition, or symptom mediated by an estrogen receptor and/or caused by lowered estrogen level in a mammal, said method comprising administering to said mammal a therapeutically effective amount of a compound of claim 1 and an amount of a second ingredient, said second ingredient being selected from the group consisting of anabolic agents, growth hormones, growth hormone secretagogues, prostaglandin agonists, prostaglandin antagonists, parathyroid hormones, sodium fluoride, prodrugs of any of the preceding, and pharmaceutically acceptable salts of any of the preceding.
10. A pharmaceutical composition comprising a compound of claim 1 and a pharmaceutically acceptable vehicle, carrier, or diluent.
11. A pharmaceutical composition comprising a compound of claim 7 and a pharmaceutically acceptable vehicle, carrier, or diluent.
12. A pharmaceutical composition comprising:
  - a compound of claim 1;
  - a second ingredient selected from the group consisting of anabolic agents, growth hormones, growth hormone secretagogues, prostaglandin agonists, prostaglandin antagonists, parathyroid hormones, sodium fluoride, prodrugs of any of the preceding, and pharmaceutically acceptable salts of any of the preceding; and
  - a pharmaceutically acceptable vehicle, carrier, or diluent.



European Patent  
Office

# PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 00 31 1197 shall be considered, for the purposes of subsequent proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	WO 97 31640 A (PFIZER; KE HUA ZHU (US); THOMPSON DAVID D (US)) 4 September 1997 (1997-09-04) * page 17, line 17 - page 22, line 9 * * claims 1-3,11,12 *	1-5, 8-10,12	C07D217/04 C07D217/06 C07D217/14 C07D217/18 C07D409/04 C07D401/12 C07D413/12
X	EP 0 792 645 A (PFIZER) 3 September 1997 (1997-09-03) * claims 1-7,12,14-17 *	1-5, 8-10,12	C07D405/06 C07D417/04 C07D401/04
P,X	WO 00 55137 A (CHAO QI; MCKIE JEFFREY; STEIN BERND; RICE KEN; GANGLOFF ANTHONY ET AL) 21 September 2000 (2000-09-21) * claims 1,16,19-21,24,25,33-44; examples 1-136 *	1-5,7-12	A61K31/472 A61K31/4725 A61P5/30
X	EP 0 792 641 A (PFIZER) 3 September 1997 (1997-09-03) * page 5, line 38 - page 6, line 18 * * page 18, line 51 - page 19, line 56 *	1-5,8,10	
X	US 5 552 412 A (CAMERON KIMBERLY O ET AL) 3 September 1996 (1996-09-03) * claims 1,8,9 *	1-5,8,10	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			C07D
-/--			
<b>INCOMPLETE SEARCH</b>			
<p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely :</p> <p>Claims searched incompletely :</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search:</p> <p>Although claims 8, 9 are directed to a method of treatment of the human/animal body (Article 52(4) EPC), the search has been carried out and based on the alleged effects of the compound/composition.</p>			
Place of search		Date of completion of the search	Examiner
BERLIN		10 May 2001	Hass, C
CATEGORY OF CITED DOCUMENTS		<p>T : theory or principle underlying the invention</p> <p>E : earlier patent document, but published on, or after the filing date</p> <p>D : document cited in the application</p> <p>L : document cited for other reasons</p> <p>&amp; : member of the same patent family, corresponding document</p>	
<p>X : particularly relevant if taken alone</p> <p>Y : particularly relevant if combined with another document of the same category</p> <p>A : technological background</p> <p>C : non-written disclosure</p> <p>P : intermediate document</p>			

EPO FORM 1503 03/82 (P04C07)



European Patent  
Office

## PARTIAL EUROPEAN SEARCH REPORT

Application Number  
EP 00 31 1197

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	EP 0 842 661 A (PFIZER) 20 May 1998 (1998-05-20) * page 2, line 5 - line 6; claims 1,4,5 *	1-5,8	
X	EP 0 792 642 A (PFIZER) 3 September 1997 (1997-09-03) * claims *	1-5,8	
X	PATENT ABSTRACTS OF JAPAN vol. 016, no. 128 (C-0924), 2 April 1992 (1992-04-02) - & JP 03 294268 A (YAMANOUCHI PHARMACEUT CO LTD), 25 December 1991 (1991-12-25) * abstract *	1,7,8, 10,11	
X	EP 0 514 851 A (SCHERING CORP) 25 November 1992 (1992-11-25) * claims 1,8,10,13 *	1,7,10, 11	TECHNICAL FIELDS SEARCHED (Int.Cl.7)
X	EP 0 286 293 A (YAMANOUCHI PHARMA CO LTD) 12 October 1988 (1988-10-12) * claims 1,5; example 11 *	1,10	
X	US 4 340 601 A (BRENNER L MARTIN) 20 July 1982 (1982-07-20) * claims 1,7 *	1,10	
X	CH 538 477 A (HOFFMANN LA ROCHE) 30 June 1973 (1973-06-30) * column 8, line 39 - line 67; examples 1,2,4-8,16,20-27; table 1 *	1,10	
X	US 3 666 763 A (GRETHE GUENTER ET AL) 30 May 1972 (1972-05-30) * abstract; examples 9,12,14-16,18,19,25,26,29-32 *	1,10	
	---		
	-/--		



European Patent  
Office

## PARTIAL EUROPEAN SEARCH REPORT

Application Number  
EP 00 31 1197

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	US 4 113 869 A (GARDNER DEREK VICTOR) 12 September 1978 (1978-09-12) * example 4, starting compound *	1	
A,D	WO 96 09040 A (LILLY CO ELI; BRYANT HENRY UHLMAN (US); CULLINAN GEORGE JOSEPH (US)) 28 March 1996 (1996-03-28) * claims 1,5,6,8,10-26 *	1-5, 8-10,12	
A,D	EP 0 826 679 A (LILLY CO ELI) 4 March 1998 (1998-03-04) * claims 1-4,11-21 *	1-5,8,10	
A,D	WO 97 13764 A (LILLY CO ELI; CULLINAN GEORGE J (US); KAUFFMAN RAYMOND F (US)) 17 April 1997 (1997-04-17) * claims 1,6-13 *	1-5,8,10	TECHNICAL FIELDS SEARCHED (Int.Cl.7)
A,D	US 5 462 950 A (FONTANA STEVEN A) 31 October 1995 (1995-10-31) * claims 1,3 *	1-5,8,10	
A	EP 0 911 321 A (PFIZER) 28 April 1999 (1999-04-28) * claims 1,17,18,22,23 *	1,8-10, 12	
A,D	US 3 277 106 A (W L BENCZE ET AL) 4 October 1966 (1966-10-04) * column 4, line 20 - line 29; claims 1,3; examples 1-3,5,6,8 *	1,8,10	

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 31 1197

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

10-05-2001

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9731640 A	04-09-1997	AU 703285 B	25-03-1999
		AU 1039897 A	16-09-1997
		BG 102726 A	30-04-1999
		BR 9612533 A	20-07-1999
		CA 2247420 A	04-09-1997
		CN 1209064 A	24-02-1999
		CZ 9802718 A	16-06-1999
		EP 0883404 A	16-12-1998
		HR 970118 A	30-04-1998
		HU 9904123 A	28-05-2000
		JP 11504352 T	20-04-1999
		NO 983936 A	27-08-1998
		PL 328831 A	15-02-1999
		SK 118398 A	11-07-2000
		ZA 9701719 A	27-08-1998
EP 0792645 A	03-09-1997	AU 1497697 A	04-09-1997
		CA 2198534 A	28-08-1997
		CN 1165654 A	26-11-1997
		JP 10007562 A	13-01-1998
WO 0055137 A	21-09-2000	AU 4172500 A	04-10-2000
EP 0792641 A	03-09-1997	AU 703473 B	25-03-1999
		AU 1497997 A	04-09-1997
		CA 2198578 A	28-08-1997
		CN 1167617 A	17-12-1997
		JP 10007563 A	13-01-1998
		US 6107331 A	22-08-2000
US 5552412 A	03-09-1996	ZA 9701714 A	27-08-1997
		AP 592 A	05-05-1997
		AU 700982 B	14-01-1999
		AU 4091696 A	18-07-1996
		BG 62256 B	30-06-1999
		BG 100278 A	31-05-1996
		BR 9600079 A	27-01-1998
		CA 2209925 A	18-07-1996
		CN 1136562 A,B	27-11-1996
		CZ 9600055 A	16-10-1996
		EP 0802910 A	29-10-1997
		FI 972903 A	08-07-1997
		HR 960010 A	31-12-1997
		HU 9600056 A	28-12-1998
		WO 9621656 A	18-07-1996
		JP 2972347 B	08-11-1999

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 31 1197

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.  
The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

10-05-2001

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5552412 A		JP 10503215 T	24-03-1998
		KR 190727 B	01-06-1999
		LV 11460 A,B	20-08-1996
		NO 960081 A	10-07-1996
		NZ 280792 A	24-11-1997
		PL 312182 A	22-07-1996
		RO 116275 B	29-12-2000
		RU 2130454 C	20-05-1999
		SG 47377 A	17-04-1998
		SI 9600004 A	31-10-1996
		SK 164895 A	07-05-1997
		TR 960693 A	21-08-1996
		US 6153622 A	28-11-2000
EP 0842661 A	20-05-1998	AU 4523797 A	21-05-1998
		HU 9702040 A	28-11-2000
		JP 10147527 A	02-06-1998
		US 6034102 A	07-03-2000
		ZA 9710292 A	17-05-1999
EP 0792642 A	03-09-1997	AU 703384 B	25-03-1999
		AU 1498097 A	04-09-1997
		CA 2198562 A	28-08-1997
		CN 1165655 A	26-11-1997
		JP 10007564 A	13-01-1998
		US 5889042 A	30-03-1999
		ZA 9701713 A	27-08-1998
JP 03294268 A	25-12-1991	NONE	
EP 0514851 A	25-11-1992	US 5124337 A	23-06-1992
		JP 7002785 A	06-01-1995
		JP 7030028 B	05-04-1995
		US 5238935 A	24-08-1993
EP 0286293 A	12-10-1988	AT 77820 T	15-07-1992
		AT 123497 T	15-06-1995
		DE 3853958 D	13-07-1995
		DE 3853958 T	05-10-1995
		DE 3872425 A	06-08-1992
		DE 3872425 T	03-12-1992
		EP 0399626 A	28-11-1990
		ES 2041790 T	01-12-1993
		ES 2075137 T	01-10-1995
		GR 3005734 T	07-06-1993
		HU 9500648 A	28-11-1995

EPO FORM P459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 31 1197

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.  
The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

10-05-2001

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0286293 A		JP 1199948 A	11-08-1989
		JP 1863983 C	08-08-1994
		JP 5075745 B	21-10-1993
		KR 9614352 B	15-10-1996
		US 4876261 A	24-10-1989
		US 5623074 A	22-04-1997
		US 4966904 A	30-10-1990
		US 5079256 A	07-01-1992
		US 5077406 A	31-12-1991
		US 5204468 A	20-04-1993
		US 5378849 A	03-01-1995
US 4340601 A	20-07-1982	AT 6061 T	15-02-1984
		DE 3162114 D	08-03-1984
		DK 300081 A,B,	09-01-1982
		EP 0043729 A	13-01-1982
		GR 74324 A	22-06-1984
		IE 51371 B	10-12-1986
		JP 57062260 A	15-04-1982
CH 538477 A	30-06-1973	CH 527194 A	31-08-1972
		AT 303041 B	15-10-1972
		BE 761219 A	05-07-1971
		CA 999866 A	16-11-1976
		DE 2062001 A	15-07-1971
		DK 125021 B	18-12-1972
		ES 387016 A	01-05-1976
		FI 49503 B	01-04-1975
		FR 2081412 A	03-12-1971
		GB 1335261 A	24-10-1973
		IE 34859 B	03-09-1975
		IL 35916 A	28-11-1973
		JP 55015476 B	23-04-1980
		NL 7018956 A,B,	08-07-1971
		NO 135315 B	13-12-1976
		NO 135062 B	25-10-1976
		NO 135314 B	13-12-1976
		PH 9682 A	10-02-1976
		SE 386672 B	16-08-1976
		SE 368401 B	01-07-1974
		SE 399879 B	06-03-1978
		US 3947456 A	30-03-1976
		YU 1971 A,B	30-06-1977
		ZA 7008590 A	29-09-1971
US 3666763 A	30-05-1972	NONE	

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82



**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 31 1197

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.  
The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

10-05-2001

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4113869 A	12-09-1978	GB 1504424 A	22-03-1978
		AT 354448 B	10-01-1979
		AT 586876 A	15-06-1979
		AT 353793 B	10-12-1979
		AT 717378 A	15-05-1979
		AU 498422 B	15-03-1979
		AU 1670276 A	16-02-1978
		BE 844783 A	31-01-1977
		CA 1076119 A	22-04-1980
		CH 626611 A	30-11-1981
		CH 622780 A	30-04-1981
		DE 2635276 A	17-02-1977
		DK 357976 A	10-02-1977
		ES 450550 A	16-12-1977
		ES 462058 A	16-06-1978
		ES 462059 A	16-06-1978
		ES 462060 A	16-06-1978
		ES 462061 A	16-06-1978
		FR 2320755 A	11-03-1977
		HU 173560 B	28-06-1979
		IE 43506 B	11-03-1981
		IL 50176 A	31-10-1979
		JP 52023083 A	21-02-1977
		NL 7608856 A	11-02-1977
		NZ 181619 A	31-08-1979
		SE 7608641 A	10-02-1977
		ZA 7604555 A	27-07-1977
WO 9609040 A	28-03-1996	AU 3727795 A	09-04-1996
		EP 0774966 A	28-05-1997
EP 0826679 A	04-03-1998	AU 721467 B	06-07-2000
		AU 4328497 A	19-03-1998
		BR 9711440 A	24-08-1999
		CN 1228758 A	15-09-1999
		CZ 9900500 A	14-07-1999
		NO 990913 A	25-02-1999
		PL 331898 A	16-08-1999
		TR 9900384 T	21-05-1999
		WO 9808797 A	05-03-1998
		US 6211184 B	03-04-2001
		ZA 9707619 A	25-02-1999
WO 9713764 A	17-04-1997	AU 7259396 A	30-04-1997
		CA 2234267 A	17-04-1997
		EP 0863892 A	16-09-1998

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 31 1197

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.  
The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

10-05-2001

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9713764	A		JP 11513664 T	24-11-1999
			US 6147092 A	14-11-2000
			US 5726168 A	10-03-1998
			US 5811415 A	22-09-1998
			US 5827844 A	27-10-1998
			US 5789400 A	04-08-1998
			US 5929092 A	27-07-1999
-----				
US 5462950	A	31-10-1995	AU 8047294 A	29-06-1995
			CA 2138100 A	22-06-1995
			CN 1108654 A	20-09-1995
			CZ 9403156 A	13-09-1995
			EP 0659421 A	28-06-1995
			HU 72313 A	29-04-1996
			IL 111995 A	31-01-2000
			JP 7215859 A	15-08-1995
			NO 944876 A	22-06-1995
			US 5770612 A	23-06-1998
			US 5760061 A	02-06-1998
			ZA 9409984 A	14-06-1996
-----				
EP 0911321	A	28-04-1999	CA 2249867 A	10-04-1999
			JP 11180926 A	06-07-1999
			US 6124314 A	26-09-2000
-----				
US 3277106	A	04-10-1966	BE 658691 A	22-07-1965
			NL 6500840 A	26-07-1965
			US 3522319 A	28-07-1970
-----				